

Abstract

Introduction

Arterial spin labeling (ASL) is a non-invasive magnetic resonance imaging (MRI) technique that can measure regional cerebral blood flow (rCBF) without radiation exposure. This study aimed to evaluate rCBF in individuals with autism and their age-matched controls, globally and regionally.

Methods

We performed ASL MRI (3T, pulsed-continuous ASL, 3 delayed ASL imaging sequences) for 33 patients with autism spectrum disorder (ASD) (average age: 7.3 years, range: 2-14 years). Nineteen children (average age: 8.6 years, range: 3-15 years) without ASD and intellectual delay were included as controls. Patients with morphological abnormalities detected on MRI were excluded. Objective analysis was performed with automatic region of interest analysis of the ASL results. The Mann-Whitney U test was used to compare the rCBF results between the groups.

Results

Compared to the controls, patients with ASD showed a statistically significant decrease in rCBF, respectively, in the insula [left, rCBF 51.8 ± 9.5 mL/100 g/min (mean \pm SD) versus 59.9 ± 9.8 , $p=0.0017$; right, 51.2 ± 10.1 versus 57.8 ± 8.8 , $p=0.0354$], superior parietal lobule (left, 44.6 ± 8.4 versus 52.0 ± 7.8 , $p=0.003$), superior temporal gyrus (left, 50.0 ± 8.6 versus 56.9 ± 8.6 , $p=0.007$; right, 49.5 ± 8.4 versus 56.4 ± 7.7 , $p=0.0058$), and inferior frontal gyrus (left, 53.0 ± 9.8 versus 59.3 ± 9.9 , $p=0.0279$), which are associated with the mirror neuron system.

Conclusions

We concluded that patients with ASD showed a statistically significant decline in CBF in regions associated with the mirror neuron system. The advantages of ASL MRI include low invasiveness (no radiation exposure) and short imaging time (approximately 5 min). Studies with larger sample sizes are required to establish the diagnostic value of ASL MRI for ASD.

Keywords: arterial spin labeling, autism spectrum disorder, cerebral blood flow,

magnetic resonance imaging, mirror neuron system

Introduction

It is thought that early detection and intervention contribute to good outcomes in social adaptation in patients with autism spectrum disorders (ASD) [1].

However, the diagnostic process depends on the physician's interview. The establishment of objective biomarkers (e.g., cerebral function, cerebral perfusion) for ASD may aid in the auxiliary diagnosis of ASD.

Functional magnetic resonance imaging (MRI) and near-infrared spectroscopy (NIRS) are useful for the examination of cerebral function [2,3]. However, these examinations involve the performance of tasks, which can be difficult for patients with developmental delay, who cannot easily follow the examiner's instructions. In contrast, regional cerebral blood flow (rCBF) single photon emission computed tomography (SPECT) does not require the performance of any tasks. Therefore, even patients with intellectual delays can undergo these investigations.

Hashimoto et al. demonstrated the decrease in frontal and temporal rCBF in the autistic brain, using CBF SPECT. The decrease in rCBF was more pronounced in patients with intellectual delay and those who were unable to use language, than in those without this disability [4]. Ohnishi et al. demonstrated that the insula, superior temporal gyrus, and prefrontal area showed a focal decrease in rCBF in young patients with ASD [5]. Further studies are needed, to establish a consensus, but the risk of radiation exposure discouraged patients with ASD and normal controls from participating in SPECT studies.

Arterial spin labeled (ASL) perfusion MRI permits noninvasive quantification of blood flow, an important physiological parameter [6]. This method involves labeling the proton spins of the inflowing water in arterial blood, by continuously inverting them at the neck region and observing the effects of this inversion on the intensity of the brain MRI. The increase in the use of 3T clinical MRI systems has dramatically improved the quality of ASL imaging, along with the development of ASL techniques. However, quantitative measurement of rCBF using ASL depends on several parameters, including T1 of brain tissue, T1 of arterial blood, and arterial transit time (ATT), which denotes the duration required by the labeled blood to travel from the labeling region to the imaged tissue. Transit time considerations are crucial in measuring absolute rCBF using ASL; thus, ASL methods using multiple post-label delay acquisitions were developed [7].

ASL is commonly used to improve the precision of 3T MRI in patients with

brain infarction and epilepsy, because of its noninvasiveness [8,9]. MRI is routinely performed in several patients with ASD, and ASL approximately requires 3 additional minutes. ASL studies for rCBF in patients with ASD have also been reported recently. Jann K et al. reported frontotemporal hyperperfusion and hypoperfusion in the dorsal anterior cingulate cortex in patients with ASD [10]. Yerys BE et al. reported hypoperfusion in the fusiform gyrus, bilaterally and in the right inferior temporal gyrus [11]. However, ASL is still being developed, and the software to analyze results objectively have not been sufficiently developed.

ASL has the potential to become a good diagnostic test for ASD. We hypothesized that a decrease in the rCBF on ASL would be observed in the regions associated with ASD. In this study, we evaluate the possibility of using ASL MRI to evaluate the biomarkers for ASD, using standardization of the brain form and automatic creation of the region of interest (ROI).

Methods

We recruited patients who were diagnosed with ASD according to DSM-V at the Tokushima University Hospital from June 2014 to September 2016. Thirty-three patients with ASD, with an average age of 7.3 years (range: 2-16 years), were recruited and evaluated using ASL MRI. We also recruited 19 children without ASD and intellectual delay, with an average age of 8.6 years (range: 3-15 years), as controls. Patients with morphological abnormalities detected on MRI were excluded from both groups. The study was approved by the Institutional Review Board and informed consent was obtained from the family members of all the children, after the purpose and risks of the study had been fully explained.

All imaging data were acquired using a 3T-MRI (Discovery 750, GE Healthcare, Waukesha, Wisconsin, U.S.A) with 16-channel head coils. Pseudocontinuous ASL, with three pulse-labeling-delays, was performed in this study. All perfusion and reference images were acquired with a 3-dimensional stack-of-spirals rapid acquisition with refocused echoes imaging sequence[12]. The imaging parameters were as follows: repetition time, 6797 ms; inversion time, 1000 ms; echo time, 11.2 ms; flip angle, 111 degrees; field of view, 240×240 mm; slice thickness, 4 mm; post labeling delay, 1.00,1.57,2.46 seconds; 6 arms with 600 samples; number of excitations, 1; and total scan

time, 5 min 36 s. Corrected CBF maps were subsequently calculated by the MRI scanner and used for evaluation.

We used the iNEUROSTAT++[®] (Nihon Medi-Physics Co Ltd, Tokyo, Japan) software to standardize the brain forms, and iSSP3.5_2tz[®] (Nihon Medi+Physics, Tokyo, Japan) to compare the ASL results between the ASD and control groups. The extracted cortical activity of the ASD group was compared to that of the control group, using a two-sample Student's *t*-test on a pixel-by-pixel basis. Calculated *t* values were converted to *Z* values, using a probability integral transformation. Following the use of iNEUROSTAT++[®] for the standardization of the brain forms, NEURO FLEXER[®] (Nihon Medi-Physics Co Ltd, Tokyo, Japan) was used for automatic ROI analysis. The analysis results were calculated by automatically segmenting the brain into 54 categories, based on the Talairach Atlas. (Figures 1 and 2)

The Mann-Whitney U test was used to compare the measured values of rCBF in each ROI of the ASD and control groups. We used the Wechsler Intelligence Scale for Children-IV to measure the intelligence quotient (IQ) scores of all participants. The correlation between measured values of rCBF and IQ scores was analyzed using linear regression. $p < 0.05$ was considered as statistically significant, for all the analyses. All data were analyzed using GraphPad Prism 6[®] (GraphPad Software Inc, La Jolla, California, U.S.A).

Results

The results of the ASL analysis with iSSP3.5_2tz[®] are as shown in Figure 3. The results of the ASL analysis with NEURO FLEXER[®] are as shown in Table 1.

Compared to the control group, patients with ASD showed a statistically significant decrease in rCBF in the subcallosal gyrus (left, $p=0.0012$; right, $p=0.0393$), insula (left, $p=0.0017$; right, $p=0.0354$), left superior parietal lobule ($p=0.003$), superior temporal gyrus (left, $p=0.007$; right, $p=0.0058$), transverse temporal gyrus (left, $p=0.0099$; right, $p=0.014$), left caudate ($p=0.0112$), left precentral gyrus ($p=0.0144$), supramarginal gyrus (left, $p=0.0389$; right, $p=0.0146$), middle temporal gyrus (left $p=0.0252$, right $p=0.0496$), uncus (left $p=0.034$, right $p=0.0255$), left inferior frontal gyrus ($p=0.0279$), right inferior temporal gyrus ($p=0.0308$), left inferior parietal lobule ($p=0.0357$), left anterior cingulate gyrus ($p=0.0374$), left postcentral gyrus ($p=0.0432$), and left lentiform

nucleus ($p=0.0474$) (Figures 4, 5, and 6), on analysis with NEURO FLEXER®.

Moreover, we further subdivided the two groups by age. Patients with ASD were subdivided into the 0-9 years subgroup [25 patients, average age :5.4 years (range: 2-9 years)] and the ≥ 10 years subgroup [8 patients, average age: 13.1 years (range: 10-16 years)]. The control group was also subdivided into the 0-9 years subgroup [11 children, 6.2 years (range: 3-9 years)] and the ≥ 10 years subgroup [8 children, 12.0 years (range: 10-15 years)]. As a result, a significant difference was discovered only in the left insula in the 0-9 years subgroup [$rCBF 50.0 \pm 7.6$ (mean \pm SD) of ASD versus 56.4 ± 11.0 in control, $p=0.0452$], and left superior parietal lobule in the ≥ 10 years subgroup (43.4 ± 7.2 versus 55.0 ± 6.3 , $p=0.0047$).

We evaluated the correlation between IQ scores and the ASL MRI results for the bilateral insulae, superior temporal gyri, and inferior frontal gyri using linear regression; however, no statistically significant correlation was observed (Figure 7).

Discussion

This study found a statistically significant reduction in the blood flow to the insula, superior temporal gyrus, left inferior frontal gyrus, and left inferior parietal lobule using ASL MRI. These areas are closely associated with the mirror neuron system and the theory of the mind, which are often impaired in children with ASD.

It has recently been proposed that dysfunction of the mirror neuron system during early development could cause a cascade of impairments that are characteristic of ASD, including deficits in imitation, theory of mind, and social communication [13]. This system is thought to provide a neural mechanism by which others' actions and intentions can be automatically understood [14].

Some human brain regions such as the lower frontal gyrus, inferior parietal lobule, and premotor area are also considered as part of the mirror-neuron system, based on functional MRI studies [15]. The visuoperceptual characteristics of the movements of others are analyzed in the superior temporal gyrus. This information is conveyed to the mirror neuron system [16]. Furthermore, this cortical system is linked to the limbic system by the anterior sector of the insular lobe [17].

Moreover, our study found a statistically significant reduction in blood flow to the subcallosal gyrus. The subcallosal gyrus sends projections to the amygdala and is involved in the suppression of the amygdala's responsiveness to fear-inducing cues. Dysfunction of this area may be responsible for the failure of extinction of the fear response, which is an important part of the anxiety response [18]. This dysfunction of the subcallosal gyrus may be related to anxiety, which several patients with ASD patients are likely to experience.

Our study also found a statistically significant reduction of blood flow to the uncus. The uncus of the hippocampus is the anterior-most portion of the medial parahippocampal gyrus and belongs to the limbic system. It is also a part of the olfactory system. It houses the primary olfactory cortex and receives fibers from the olfactory tract via the lateral olfactory stria [19]. Decreased blood flow to the uncus may be related to the olfactory and limbic abnormalities in ASD.

Furthermore, our study found a statistically significant reduction of blood flow to the transverse temporal gyrus. This structure (also called Heschl's gyrus) is a part of the primary auditory cortex [20]. Reduced blood flow to this region may be correlated with difficulties that autistic children experience, while processing auditory stimuli, compared to visual stimuli.

We also analyzed the rCBF in segmentalized age-groups. A significant difference was observed only in the left insula in the 0-9 years group and the left superior parietal lobule in the ≥ 10 years group. This result reflects the fact that it becomes difficult to determine a significant difference if the size of the sample is too small. However, these regions are associated with the mirror neuron system, thus reinforcing our results.

Finally, the lack of correlation between the IQ scores and measured decrease in rCBF suggests that the difficulty experienced by patients with ASD in daily life is not necessarily correlated with their IQ.

The present study has some limitations. First, some children in the control group showed reduction of rCBF in the ROI related to the mirror neuron system. Therefore, we were unable to diagnose ASD by ASL MRI alone. Correlation of the results of diffusor tensor imaging/functional MRI with rCBF results obtained by ASL could have aided the diagnosis. Second, an autism assessment tool such as the Parent-interview ASD Rating Scale-Text Revision should have been used, to better evaluate the relationship between ASD characteristics and rCBF. Third, the site of the hypoperfusion reported by

previous ASL study in patients with ASD [9,10] were different from our study. However, several points are similar to the SPECT study, including hypoperfusion in the regions associated with the mirror neuron system [4,5].

Conclusion

In this study, ASL MRI demonstrated a statistically significant decline in rCBF in regions associated with the mirror neuron system in patients with ASD. ASL MRI had potential as a useful diagnostic technique for ASD. ASL MRI is highly advantageous, owing to its noninvasiveness (no radiation exposure) and short imaging time (approximately 5 min). However, the sample size of our study was small. Further studies with larger sample sizes are required, to irrefutably establish the diagnostic value of ASL MRI. It is also necessary to evaluate the correlation between the severity of ASD and the reduction in rCBF measured by ASL MRI, using autism assessment scales such as the Parent-interview ASD Rating Scale-Text Revision.

Declaration of Competing Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

References

1. Dawson G, Rogers S, Munson J, Smith M, Winter J, Greenson J, et al. Randomized, controlled trial of an intervention for toddlers with autism: the Early Start Denver Model. *Pediatrics* 2010;125:e17–23.
2. Wang AT, Lee SS, Sigman M, Dapretto M. Reading Affect in the Face and Voice: Neural Correlates of Interpreting Communicative Intent in Children and Adolescents with Autism Spectrum Disorders. *Arch Gen Psychiatry* 2007;64:698–708.
3. Mori K, Toda Y, Ito H, Mori T, Mori K, Goji A, et al. Neuroimaging in autism spectrum disorders: 1H-MRS and NIRS study. *J Med Invest* 2015;62:29–36.
4. Hashimoto T, Sasaki M, Fukumizu M, Hanaoka S, Sugai K, Matsuda H. Single-photon emission computed tomography of the brain in autism: effect of the developmental level. *Pediatr Neurol* 2000;23:416–20.
5. Ohnishi T, Matsuda H, Hashimoto T, Kunihiro T, Nishikawa M, Uema T, et al. Abnormal regional cerebral blood flow in childhood autism. *Brain* 2000;123:1838–44.
6. Alsop DC, Detre JA, Golay X, Günther M, Hendrikse J, Hernandez-Garcia L, et al. Recommended implementation of arterial spin-labeled perfusion MRI for clinical applications: A consensus of the ISMRM perfusion study group and the European consortium for ASL in dementia. *Magn Reson Med* 2015;73:102-16.
7. Tsujikawa T, Kimura H, Matsuda T, Fujiwara Y, Isozaki M, Kikuta K, et al. Arterial Transit Time Mapping Obtained by Pulsed Continuous 3D ASL Imaging with Multiple Post-Label Delay Acquisitions: Comparative Study with PET-CBF in Patients with Chronic Occlusive Cerebrovascular Disease. *PLoS One* 2016;11:e0159894.
8. Okazaki S, Yamagami H, Yoshimoto T, Morita Y, Yamamoto H, Toyoda K, et al. Cerebral hyperperfusion on arterial spin labeling MRI after reperfusion therapy is related to hemorrhagic transformation. *J Cereb Blood Flow Metab* 2017;37:3087–90.
9. Ho ML. Arterial spin labeling: Clinical applications. *J Neuroradiol* 2018;45:276–89.
10. Jann K, Hernandez LM, Beck-Pancer D, McCarron R, Smith RX, Dapretto M, et al. Altered resting perfusion and functional connectivity of default mode network in youth with autism spectrum disorder. *Brain Behav* 2015;5:e00358.

11. Yerys BE, Herrington JD, Bartley GK, Liu HS, Detre JA, Schultz RT. Arterial spin labeling provides a reliable neurobiological marker of autism spectrum disorder. *J Neurodev Disord* 2018;10:32.
12. Dai W, Shankaranarayanan A, Alsop DC. Volumetric measurement of perfusion and arterial transit delay using hadamard encoded continuous arterial spin labeling. *Magn Reson Med*. 2013;69:1014-22.
13. Dapretto M, Davies MS, Pfeifer JH, Scott AA, Sigman M, Bookheimer SY, et al. Understanding emotions in others: mirror neuron dysfunction in children with autism spectrum disorders. *Nat Neurosci* 2006;9:28–30.
14. Rizzolatti G, Craighero L. The mirror-neuron system. *Annu Rev Neurosci* 2004;27:169–92.
15. Iacoboni M, Woods RP, Brass M, Bekkering H, Mazziotta JC, Rizzolatti G. Cortical mechanisms of human imitation. *Science* 1999;286:2526–8.
16. Binkofski F, Buccino G, Posse S, Seitz RJ, Rizzolatti G, Freund H. A fronto-parietal circuit for object manipulation in man: evidence from an fMRI-study. *Eur J Neurosci* 1999 11:3276–86.
17. Hoffman M. How automatic and representational is empathy, and why. *Behav Brain Sci* 2002;25:38-9.
18. Vermetten E, Lanius RA. Biological and clinical framework for posttraumatic stress disorder. *Handb Clin Neurol* 2012;106:291–342.
19. Radiopaedia.org[Internet]. Uncus. [cited 2019 Jun 5]. Available from: <https://radiopaedia.org/articles/uncus>.
20. Amunts K, Morosan P, Hilbig H, Zilles K. Auditory System. In: Juergen Mai, George Paxinos, editors. *The Human Nervous System (Third Edition)* Cambridge: Academic Press Inc; 2012. p.1270-94.

Figure legends

Figure 1

Regions of interest (ROI) as seen on Neuro Flexer®.

The results of the analysis can be calculated, by automatically segmenting the brain into 54 categories, based on the Talairach Atlas, with the Level 3 setting of this software. The numbers are the same as in Figure 2.

Figure 2

Results of Neuro Flexer® analysis for an 8-year-old boy in the control group and an 8-year-old boy in the ASD group.

ASD: autism spectrum disorder

Figure 3

Comparison of the ASL results between the ASD and control groups, as seen on iSSP3.5_2tz®.

ASL: arterial spin labeling. ASD: autism spectrum disorder

Figure 4

Statistically significant blood flow reduction detected by arterial spin labeling magnetic resonance imaging of the insula (left $p=0.0017$), superior temporal gyrus (left $p=0.007$), inferior frontal gyrus (left $p=0.028$), and inferior parietal lobule (left $p=0.036$) These areas are closely associated with the mirror neuron system and the theory of the mind, which are often impaired in children with autism spectrum disorders. These regions are essential for sympathy and for understanding others' intentions.

Figure 5

Statistically significant blood flow reduction detected by arterial spin labeling magnetic resonance imaging in the subcallosal gyrus (left, $p=0.0012$; right, $p=0.039$) The subcallosal gyrus sends projections to the amygdala and is involved in the suppression of the amygdala's responsiveness to fear-inducing cues.

Figure 6

Statistically significant blood flow reduction detected by arterial spin labeling magnetic resonance imaging in the uncus (left, $p=0.034$; right, $p=0.026$) The uncus of the hippocampus is a part of the limbic and olfactory systems.

Decreased blood flow to the uncus of the hippocampus may be related to the olfactory and limbic abnormalities seen in ASD.

ASD: autism spectrum disorder

Figure 7

Linear regression analysis showing a lack of statistically significant correlation between IQ scores and arterial spin labeling magnetic resonance imaging results for the bilateral insulae, superior temporal gyri, and inferior frontal gyri

IQ: intelligence quotient