

Protective effects of resveratrol against 5-fluorouracil-induced oxidative stress and inflammatory responses in human keratinocytes

Shu Chen,¹ Naofumi Tamaki,^{1,*} Yasusei Kudo,² Takaaki Tsunematsu,³ Kaname Miki,¹ Naozumi Ishimaru,³ and Hiro-O Ito¹

¹Department of Preventive Dentistry, ²Department of Oral Bioscience, and ³Department of Oral Molecular Pathology, Tokushima University Graduate School of Biomedical Sciences, 3-18-15 Kuramoto-cho, Tokushima 770-8504, Japan

(Received 19 February, 2021; Accepted 11 June, 2021; Published online 2 October, 2021)

Although 5-fluorouracil (5-FU) is currently used as an anti-cancer chemotherapy, adverse effects such as oral mucositis potentially limit its clinical application. Additionally, the prevention of 5-FU-induced side effects are scarce. Resveratrol is known to decrease oxidative damage and inflammation. In this study, we examined the protective effects of resveratrol on 5-FU-induced oxidative stress and inflammatory responses in normal human keratinocytes (HaCaT cell) as *in vitro* oral mucositis model. HaCaT cells were exposed to 5-FU and simultaneously treated with resveratrol. The effects of resveratrol on 5-FU-induced cytotoxicity were evaluated using cell viability assay. The production of reactive oxygen species (ROS) was measured using a fluorescence spectrophotometer. The effects of resveratrol on nuclear factor erythroid 2-related factor 2 (Nrf2), silent information regulator transcript-1 (SIRT-1), and nuclear factor kappa B (NF-κB) signaling and inflammatory cytokine expression were examined. Resveratrol suppressed 5-FU-induced overproduction of ROS by upregulating anti-oxidant defense genes through Nrf2 activation and SIRT-1 expression. Concerning inflammatory responses, resveratrol suppressed the 5-FU-induced expression of pro-inflammatory cytokines via NF-κB nuclear translocation. Conversely, *N*-acetylcysteine reduced ROS levels without affecting the expression of pro-inflammatory cytokines. Resveratrol might be useful for preventing 5-FU-induced adverse effects by activating anti-oxidant and anti-inflammatory responses.

Key Words: oxidative stress, inflammation, oral mucositis, resveratrol, human normal keratinocyte

The pyrimidine anti-metabolite 5-fluorouracil (5-FU) is widely used to treat a variety of cancers; however, it causes several side effects. Oral mucositis, a painful inflammatory ulcer of the oral mucosa, is a common side effect occurring in up to 90% in patients with cancer treated with 5-FU.^(1,2) The clinical manifestation of oral mucositis is ulceration of the mucosal epithelium. In addition, the formation of ulcers causes severe pain, altered taste, dry mouth, and decreased appetite, and it directly or indirectly induces the suppression of nutritional uptake in many cases.⁽³⁻⁵⁾ Despite the availability of palliative treatments including antibiotics, analgesics, and radioprotective agents, in addition to general oral hygiene modification, these symptomatic therapies have limited efficacy.⁽⁶⁾

Recent studies illustrated that the pathogenesis of oral mucositis is based on reactive oxygen species (ROS) generation and inflammation.^(2,3) The overproduction of ROS leads to their accumulation and oxidative stress, which further aggravate cell damage and eventually cause apoptosis.⁽⁷⁾ Therefore, cells

scavenge the overproduced ROS through anti-oxidant networks, and the balance between ROS production and scavenging is tightly regulated.⁽⁸⁾ Nuclear factor erythroid 2-related factor 2 (Nrf2) is a pleiotropic protein and important anti-oxidant sensor. Under oxidative stress, electrophiles and oxidants modify Kelch-like ECH-associated protein-1, leading to its dissociation from Nrf2. Following its release, Nrf2 translocates to the nucleus and then binds to an anti-oxidant response element (ARE) sequence to mediate the transcription of target genes, such as heme oxygenase 1 (HO-1) and NAD(P)H-quinone oxidoreductase (NQO-1).⁽⁹⁾ Transcription of these genes increases the resistance of cells to oxidative stress and protects against inflammation.⁽¹⁰⁾

Resveratrol is a polyphenolic phytoalexin that naturally occurs in many plant products, such as grapes, berries, and peanuts.⁽¹¹⁾ It has been reported that resveratrol is more effective than conventional anti-oxidants such as *N*-acetylcysteine (NAC) and quercetin in decreasing oxidative damage.⁽¹²⁾ The cellular effects of resveratrol were attributed to its anti-inflammatory and anti-oxidant capacities. However, the protective effects of resveratrol on 5-FU-induced damage in normal keratinocytes remain to be determined. Furthermore, resveratrol has been reported to enhance 5-FU-induced cytotoxicity in cancer cells.⁽¹³⁾

In this study, a normal human keratinocyte cell line (HaCaT) was used as an *in vitro* model to investigate oral mucositis caused by 5-FU, as these cells can serve as a useful model for oral mucositis research.⁽¹⁴⁾ Several studies suggested that HaCaT cells can produce pro-inflammatory cytokines, such as interleukin (IL)-1, IL-6,⁽¹⁵⁾ and tumor necrosis factor (TNF)-α.⁽¹⁶⁾ However, the roles of ROS production and upregulation of the nuclear factor kappa B (NF-κB) pathway induced by 5-FU in normal keratinocytes have not been explored.

We hypothesized that resveratrol can prevent or attenuate oral mucositis caused by 5-FU. We evaluated the inhibitory effects of resveratrol on 5-FU-induced oxidative stress and inflammatory responses. Moreover, we investigated the effects of resveratrol on the Nrf2, silent information regulator transcript-1 (SIRT-1), and NF-κB signaling pathways.

Materials and Methods

Cell culture. The HaCaT immortalized keratinocyte cell line from adult human skin was obtained from German Cancer Research Center (Heidelberg, Germany). Cells were plated in cell culture flasks and suspended in Dulbecco's Modified Eagle

*To whom correspondence should be addressed.
E-mail: tamanao@tokushima-u.ac.jp

Table 1. Primers used in this study for real-time PCR

Gene Name	Forward Primer	Reverse Primer
IL-1 β	TGATGGCTTATTACAGTGGC	TGTAGTGGTGGTCGGAGATT
IL-6	AAGCCAGAGCTGTCAGATGAGTA	TGTCTGCAGCCACTGGTTC
TNF- α	CCTCTGGCCCAGGCAGTCAGA	GGCGTTGGGAAGGTTGGAT
IL-10	GCTGCACCCACTCCAGGC	CCTGCTCACGGCCTTGCTC
HO-1	CTTCAGAAGGGTCAGGTGTC	TGCTTCTTCGCTCTATCTCC
NQO-1	CATCATTTGGGTCAGTCC	ACAGCCGTGGCAGAACTA
SIRT-1	AAGGAAAACACTTCGCAAC	GGAACCATGACACTGAATTA
GAPDH	GTATTGGGCCTGGTACCC	CGCTCCTGGAAGAGATGGTGATGG

Medium (DMEM; Nacalai Tesque, Inc., Kyoto, Japan) supplemented with 10% fetal bovine serum (MP Biomedicals LLC, Santa Ana, CA), 100 U/ml penicillin, 0.1 mg/ml streptomycin (Sigma-Aldrich, St. Louis, MO), and kept in a humidified incubator at 37°C, with an atmosphere of 5% CO₂.

Cell viability. The cell viability assay was conducted to determine the concentrations of resveratrol as a chemopreventive agent against toxicity induced by 5-FU. Briefly, HaCaT cells were seeded into the 96-well plate and incubated for 24 h. Then cells treated with or without different concentrations of resveratrol and 5-FU for 24 h. WST-1 (Takara Bio Inc., Otsu, Japan) assay was performed according to the standard protocol of manufacture. Absorbance value was measured at 450 nm by a microplate reader. Cell viability was expressed as the percentage of untreated cells (control).

Production and detection of ROS. HaCaT cells were incubated with 5-FU in presence or absence of resveratrol for 24 and 48 h. The adherent cells were stained with CellROX Green Reagent (Life Technology, Carlsbad, CA) at 37°C in dark for 30 min, nuclei were stained by 4',6-diamidino-2-phenylindole (DAPI) for 15 min, and harvested in PBS. The cell suspension (10,000 cells/well) was placed in a flat bottom 96-well assay plate (Nunc/ThermoFisher scientific, Waltham, MA). The intensity was then immediately measured using the fluorescence spectrophotometer (Tecan M200 Infinite PRO, Tecan Japan Co, Kawasaki, Japan) at an excitation wavelength 485 nm and emission wavelength 520 nm.

Immunofluorescence staining. HaCaT cells were incubated and then treated with 5-FU in the presence or absence of resveratrol for 24 h. Subsequently, the cells were fixed with 4% paraformaldehyde for 15 min followed by permeabilized with 0.5% Triton X-100. The cells were blocked with Blocking One Histo (Nacalai Tesque) for 10 min to prevent non-specific labelling. The cells were incubated with a monoclonal antibody specific to Nrf2 (sc-722, Santa Cruz Biotechnology, Dallas, TX) overnight at 4°C, then incubated with a secondary antibody, anti-rabbit IgG Fab2 Alexa Fluor 488 conjugate (Cell Signaling Technology, Inc., Beverly, MA) for 1 h. Nuclei were stained with DAPI for 15 min. Images were captured using a fluorescent microscopy BZ-X800 (Keyence Co., Osaka, Japan) at 400× magnification.

Extraction of protein and Western blot analysis. The nuclear protein extracts were prepared from HaCaT cells using NE-PER Nuclear and Cytoplasmic Extraction Kit (Thermo Scientific, Rockford, IL), according to the manufacturer's protocol. Additionally, total protein was extracted by ice-cold RIPA buffer (Nacalai Tesque) with protease inhibitor cocktail. The protein samples were subsequently prepared 4× Laemmli Sample Buffer (Bio-Rad Laboratories, Forester City, CA) and boiled at 100°C for 5 min. Equal amounts of protein were separated by SDS-PAGE and then transferred onto polyvinylidene difluoride membranes for 35 min at 100 V. After that, the membranes were blocked with Blocking One (Nacalai Tesque)

for 40 min at room temperature. The membranes were incubated at 4°C overnight in each primary antibody. Primary antibodies specific to SIRT-1, HO-1, NF-κB, histone H3 and β-actin (Cell Signaling Technology) and Nrf2 were used. Membranes were washed by TBST three times and incubated with anti-rabbit or anti-mouse horseradish peroxidase-conjugated secondary antibody (Cell Signaling Technology). To detect the immunocomplex, an Immobilon HRP substrate (Bio-Rad Laboratories) was used. Immunoblots were scanned by densitometry and the intensity was quantified by using Image Lab software (Bio-Rad Laboratories). The relative levels of SIRT-1 and HO-1 were normalized to β-actin, and nuclear Nrf2 and NF-κB were standardized to histone H3.

Real-time PCR. HaCaT cells were sub-cultured and treated with 5-FU in the presence or absence of resveratrol and NAC (Sigma-Aldrich) for 24 and 48 h. Total RNA was extracted using Trizol reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. Total RNA was reverse transcribed using ReverTra Ace (Toyobo, Osaka, Japan). Real-time PCR was performed using SYBR Green (Bio-Rad Laboratories) and a real-time PCR system (Bio-Rad Laboratories) according to the following cycling parameters: initial denaturation at 95°C for 1 min, 40 cycles of denaturation at 95°C for 20 s, annealing at 60°C for 30 s and extension at 72°C for 25 s. The primer sequences were showed in Table 1. The mRNA levels were calculated by determining the relative expression compared with GAPDH and the Ct of the control was normalized to 1.

ELISA assay. HaCaT cells were seeded and preincubated for 24 h. For the next 48 h, the cells treated with 5-FU in the presence or absence of resveratrol. The culture supernatants were collected from both of treated and untreated cells, and centrifuged to remove cell debris. We used commercially available ELISA kits (Biologen, San Diego, CA), according to the manufacturer's instructions. The concentrations of cytokines were calculated according to standard curves, and the obtained results are shown as the amount of IL-1β, IL-6, and TNF-α (pg/ml).

Statistical analysis. For statistical analysis, differences between the groups were analyzed with one-way analysis of variance (ANOVA) and Tukey's post hoc test for multiple comparisons using JMP (ver. 12, SAS Institute, Cary, NC). A value of $p < 0.05$ was regarded as statistically significant.

Results

Effects of resveratrol on 5-FU-induced cytotoxicity in HaCaT cells. The cytotoxicity of resveratrol and 5-FU (5 μg/ml) in HaCaT cells was evaluated after 24 h of treatment. We first examined the effects of various concentrations of resveratrol (0.1–1,000 μM) on the viability of HaCaT cells in the presence and absence of 5-FU. Higher concentrations of resveratrol (>100 μM) reduced HaCaT cell viability (Fig. 1A), whereas lower concentrations (≤ 100 μM) had no influence on




Fig. 1. Effects of resveratrol on 5-FU-induced cytotoxicity in HaCaT cells. HaCaT cells were treated with different concentrations of resveratrol alone (A) or 5-FU (5 µg/ml) and resveratrol (B) for 24 h, and then cell viabilities were evaluated by WST-1 assay. The cell viabilities were expressed as % of control. Results are presented as the means ± SD ($n = 6$). ** $p < 0.01$, *** $p < 0.001$ significantly different from control. †† $p < 0.01$, ††† $p < 0.001$ significantly different from 5-FU alone.




Fig. 2. Effects of resveratrol on the ROS production in HaCaT cells treated with 5-FU. HaCaT cells were incubated with 5-FU in the presence or absence of resveratrol for 24 h (A) and 48 h (B). Results are presented as the means ± SD of three independent experiments. ** $p < 0.01$, *** $p < 0.001$ significantly different from control. †† $p < 0.001$ significantly different from 5-FU alone.

viability. In addition, 5-FU (5 µg/ml) treatment alone for 24 h significantly decreased cell viability relative to the control ($62.7 \pm 0.3\%$). Although not all the lower concentrations of resveratrol had significantly increased the viability compared with 5-FU alone, there was a tendency for lower concentrations (1–100 µM) to protect cytotoxicity induced by 5-FU (Fig. 1B).

Resveratrol suppresses 5-FU-induced ROS generation by activating the Nrf2 pathway in HaCaT cells. To investigate the anti-oxidant effects of resveratrol, we measured ROS production in HaCaT cells. 5-FU treatment for 24 or 48 h resulted in significantly increased ROS production (Fig. 2). Additionally, resveratrol significantly decreased ROS production induced by 5-FU treatment (Fig. 2).

We examined the effects of resveratrol on the Nrf2/anti-oxidant pathway. Nuclear expression of Nrf2 was slightly observed in 5-FU-treated HaCaT cells, but resveratrol increased nuclear Nrf2 expression as observed via immunofluorescence and western blot analyses using nuclear extracts (Fig. 3A and B). Thus, resveratrol accelerated Nrf2 nuclear translocation in HaCaT cells. Addition-

ally, the expression of the anti-oxidant enzymes NQO-1 and HO-1 were analyzed by real-time PCR. 5-FU and resveratrol increased the expression of NQO-1 and HO-1 compared to their control levels (Fig. 3D and F). We also found that the protein expression of HO-1 was significantly increased by resveratrol treatment compared to the effects of 5-FU alone (Fig. 3C). Moreover, resveratrol alone also increased the expression of NQO-1 and HO-1 compared to control levels (Fig. 3E and G). These findings indicate that resveratrol activates the Nrf2 pathway and upregulates anti-oxidant defense genes.

Resveratrol increases the levels of SIRT-1 in HaCaT cells. Resveratrol is known as a SIRT-1 activator. Indeed, the mRNA expression of SIRT-1 was increased by resveratrol compared to control level in HaCaT cells (Fig. 4A). Additionally, we examined SIRT-1 expression after 5-FU treatment with or without resveratrol using real-time PCR and western blotting. 5-FU treatment increased the mRNA expression of SIRT-1 in HaCaT cells (Fig. 4B). Moreover, resveratrol synergistically enhanced both the protein and mRNA levels of SIRT-1 compared




Fig. 3. Effects of resveratrol on the Nrf2 nuclear translocation, and expressions of HO-1 and NQO-1 in HaCaT cells treated with 5-FU. HaCaT cells incubated with 5-FU in the absence or presence of resveratrol for 24 h. (A) Immunofluorescence staining of Nrf2 for each treatment. Nuclei were labeled with DAPI. 400x magnification. (B, C) Representative blots and quantitative analyses are shown for nuclear Nrf2, histone H3, HO-1, and β-actin. (D–G) Relative mRNA levels of NQO-1 and HO-1 were determined by real-time PCR. Bars represent mRNA expression normalized to GAPDH and relative to the control. Results are presented as the means \pm SD ($n = 6$). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ significantly different from control. # $p < 0.05$, ## $p < 0.01$ significantly different from 5-FU alone. ### $p < 0.001$ significantly different from resveratrol (10 µM) alone.




Fig. 4. Effects of resveratrol on the levels of SIRT-1 in HaCaT cells treated with 5-FU. HaCaT cells incubated with 5-FU in the absence or presence of resveratrol for 24 h. (A, B) Relative mRNA levels of SIRT-1 were determined by real-time PCR. Bars represent mRNA expression normalized to GAPDH and relative to the control. (C) Representative blots are shown for SIRT-1 and β -actin. Quantitative analyses are shown for SIRT-1/ β -actin. * $p<0.05$, ** $p<0.01$, *** $p<0.001$ significantly different from control. † $p<0.05$, ‡ $p<0.01$, †† $p<0.001$ significantly different from 5-FU alone. # $p<0.05$ significantly different from resveratrol (10 μM) alone.

to the effects of 5-FU alone (Fig. 4B and C).

Resveratrol inhibits 5-FU-induced NF- κ B p65 nuclear translocation in HaCaT cells. The effect of resveratrol on the 5-FU-induced nuclear translocation of the NF- κ B p65 subunit was examined (Fig. 5A). NF- κ B p65 expression in the nucleus was significantly increased by 5-FU treatment, and resveratrol reduced the 5-FU-induced nuclear expression of p65. It is thus suggested that resveratrol inhibits the NF- κ B pathway, which is activated by 5-FU. We also examined the effect of resveratrol alone on NF- κ B expression. Resveratrol treatment had no significant influence on its expression (Fig. 5B).

Resveratrol suppresses 5-FU-induced inflammatory cytokine expression in HaCaT cells. The mRNA and protein levels of inflammatory cytokines in HaCaT cells were measured by real-time PCR (Fig. 6A–H) and ELISA (Fig. 6I and J), respectively. Significantly higher expression of IL-1 β , IL-6, and TNF- α was observed after exposure to 5-FU at different time points (24 and 48 h; Fig. 6A–C and 6E–G, respectively). Resveratrol significantly decreased the expression of IL-1 β , IL-6, and TNF- α induced by 48 h of 5-FU treatment (Fig. 6E–G). Further, the protein levels of IL-1 β and IL-6, which were increased by 5-FU treatment, were also reduced after 48 h of resveratrol treatment (Fig. 6I–J). TNF- α protein levels were under the detection limit of ELISA in all conditions. We also investigated the effect of resveratrol on IL-10 expression, finding that the treatment had no significant influence on its expression (Fig. 6D, H, and K).

NAC suppresses 5-FU-induced ROS production but not pro-inflammatory cytokine expression. We examined whether NAC could decrease the expression of pro-inflammatory cytokines by suppressing the overproduction of ROS. As expected, NAC significantly decreased ROS production induced by 5-FU treatment (Fig. 7A). However, NAC did not reduce the expression of inflammatory cytokines induced by 5-FU treatment at 24 (Fig. 7B–D) and 48 h (Fig. 7E–G). This finding suggests that the expression of inflammatory cytokines in HaCaT cells

following 5-FU treatment is not a downstream event of ROS production and is independent of oxidative stress.

Discussion

In this study, we explored the utility of resveratrol for treating chemotherapy-induced oral mucositis using an *in vitro* cell culture model. To the best of our knowledge, this was the first study to demonstrate the protective effects of resveratrol on oxidative stress and inflammation induced by chemotherapy in normal human keratinocytes. Additionally, our findings suggest that resveratrol exerts substantial anti-oxidant and anti-inflammatory effects by activating Nrf2 signaling and inhibiting NF- κ B signaling, respectively.

5-FU has excellent effects against a variety of cancers including breast, head and neck, stomach, and skin cancers.⁽²⁾ 5-FU blocks DNA replication and induces the overproduction of ROS and upregulation of inflammatory responses, thereby directly inducing toxicity in the oral mucosa.^(17–19) 5-FU commonly causes oral mucositis in patients with cancer.⁽¹⁾ The pathogenesis of oral mucositis is divided into five biological steps: (i) initiation, (ii) primary damage response, (iii) signal amplification, (iv) ulceration, and (v) healing.⁽⁴⁾ During the initial stage of oral mucositis, anti-cancer drugs induce the overproduction of ROS, which play a primary role in the activation of a variety of transcription factors, such as NF- κ B.⁽²⁰⁾ Activation of NF- κ B subsequently leads to the production of the inflammatory cytokines IL-1 β , IL-6, and TNF- α .⁽¹⁹⁾ The inflammatory cytokines interact synergistically with NF- κ B, thereby amplifying the initial signals toward cell damage and tissue destruction, which occur in the primary damage response of oral mucositis.^(20,21) We revealed that the nuclear translocation of NF- κ B p65 (Fig. 5) and production of pro-inflammatory cytokines (Fig. 6) were significantly induced by 5-FU treatment in HaCaT cells, suggesting that the nuclear translocation of NF- κ B




Fig. 5. Effects of resveratrol on nuclear translocation of NF-κB in HaCaT cells treated with 5-FU. (A) HaCaT cells were pre-treated with resveratrol for 3 h and then added 5 μg/ml of 5-FU for 6 h. Representative Western blots and quantitative analysis were shown. (B) Relative mRNA levels of NF-κB were determined by real-time PCR. Bars represent mRNA expression normalized to GAPDH and relative to the control. ** $p<0.01$ significantly different from control. † $p<0.05$, †† $p<0.01$ significantly different from 5-FU alone.

leads to the upregulation of pro-inflammatory cytokines.

5-FU treatment increased the levels of ROS in HaCaT cells (Fig. 2). There is a balance between the production and elimination of ROS in the healthy condition. However, chronic inflammation leads to sustained ROS overproduction. Therefore, it is believed that the uptake of exogenous anti-oxidants is beneficial for eliminating ROS and controlling oxidative stress in the body.^(22,23) Previous researches have illustrated that resveratrol is the efficient hydroxyl radical and superoxide radical scavenger, which is substantial but less effective than known radical scavengers such as glutathione and ascorbic acid.^(22,24) In addition, resveratrol has been reported to increase the levels of anti-oxidant genes, i.e., Nrf2, HO-1, and SIRT-1, in normal cells.^(25,26) It has been also reported that resveratrol can regulate the expression and activity of anti-oxidant enzymes via the activation of the transcription factor Nrf2.⁽²⁷⁾ Our findings demonstrated that resveratrol could reduce the overproduction of ROS induced by 5-FU in HaCaT cells (Fig. 2). Moreover, resveratrol can promote the activity of SIRT-1, which regulates the activity of multiple downstream targets including the NF-κB pathway.^(28,29) Consistent with these previous studies, resveratrol could control oxidative stress with activating of Nrf2/anti-oxidant and SIRT-1 defense pathways as well as ROS scavenging.

Nrf2 is considered a main regulator of anti-oxidant defenses in

mammals.⁽³⁰⁾ In response to oxidative stress, Nrf2 is activated, and it translocates to the nucleus, in which it binds to AREs and upregulates the expression of phase 2 enzymes, such as HO-1 and NQO-1, thereby maintaining the redox balance of cells.^(31,32) We found that the nuclear translocation of Nrf2 was more strongly increased in HaCaT cells treated with both 5-FU and resveratrol than in cells treated with 5-FU alone (Fig. 3). In addition, the upregulation of HO-1 and NQO-1 expression was significantly induced by resveratrol (Fig. 3). As 5-FU-induced ROS overproduction was significantly suppressed by resveratrol (Fig. 2), the compound might regulate oxidative stress by activating the Nrf2/anti-oxidant pathway in normal cells.

SIRT-1 is a nicotinamide adenine dinucleotide-dependent nuclear histone deacetylase that participates in the regulation of several processes, such as inflammation, histone deacetylation, and apoptosis.^(33–35) SIRT-1 reduces the levels of oxidative stress and the severity of inflammation.^(36,37) Moreover, SIRT-1 promotes the nuclear accumulation, DNA binding, and transcriptional activities of Nrf2 by upregulating its downstream genes HO-1 and NQO-1.⁽³⁸⁾ In addition, the inhibition of Nrf2 ubiquitination by SIRT-1, which increases the stability and activity of Nrf2, suppresses apoptosis and resistance to oxidative stress.⁽³⁹⁾ In this study, we illustrated that resveratrol enhanced both SIRT-1 and nuclear Nrf2 levels in HaCaT cells (Fig. 3 and 4), suggesting that SIRT-1 promotes the stability and activity of Nrf2 under treatment with resveratrol. SIRT-1 also inhibits the transactivation of NF-κB.⁽³⁵⁾ In addition, HO-1 has a central role in NF-κB inhibition in the cytoplasm.⁽⁴⁰⁾ Therefore, we speculate that resveratrol can suppress the inflammatory responses activated by NF-κB through upregulating SIRT-1 and HO-1.

NF-κB activation is a cellular injury response that leads to inflammation via the production of pro-inflammatory cytokines, such as IL-1β, IL-6, and TNF-α.⁽⁴¹⁾ In an inactive state, NF-κB exists in the cytoplasm as a heterodimer of p65 and p50 subunits, but once activated by the phosphorylation and degradation of inhibitor of kappa B, it translocates to the nucleus and activates the transcription of various pro-inflammatory genes.⁽⁴²⁾ Resveratrol does not affect the DNA-binding activity of NF-κB, but it inhibits the nuclear translocation of p65.⁽⁴³⁾ In the present study, differing from resveratrol, NAC significantly decreased the overproduction of ROS induced by 5-FU, and it did not affect the expression of inflammatory cytokines (Fig. 7). This suggests that 5-FU directly induces the upregulation of inflammatory cytokines rather than through the overproduction of ROS. Importantly, resveratrol might protect cells against the cytotoxicity of 5-FU via two independent pathways, i.e., anti-oxidant and anti-inflammatory signaling. In the latter, suppression of the 5-FU-induced nuclear translocation of NF-κB p65 is likely responsible for the downregulation of inflammatory cytokines.

In conclusion, resveratrol displayed multiple beneficial effects in an *in vitro* oral mucositis model induced by 5-FU. Resveratrol decreased 5-FU-induced ROS production by activating the Nrf2 and SIRT-1 pathways in normal human keratinocytes. Moreover, resveratrol suppressed inflammatory responses by reducing the production of inflammatory cytokines via the suppression of NF-κB signaling. Overall, our results support the hypothesis that resveratrol is a safe and beneficial candidate for preventing oral mucositis associated with cancer chemotherapy. However, further studies are required before the practical use of resveratrol supplements to prevent oral mucositis.

Author Contributions

SC contributed to perform all experiments and drafted the manuscript. NT contributed to design the study, perform experiments, and revise manuscript. YK contributed to perform immunofluorescence staining, and revise manuscript. TT and KM contributed to perform Western blot analysis. NI and HI




Fig. 6. Effects of resveratrol on 5-FU induced-inflammatory cytokines in HaCaT cells treated with 5-FU. HaCaT cells were treated with 5-FU in the presence or absence of resveratrol for 24 h (A–D, K) and 48 h (E–H). IL-1 β , IL-6, TNF- α , and IL-10 levels were determined by real-time PCR. Bars represent mRNA expression normalized to GAPDH and relative to control. (I) IL-1 β and (J) IL-6 amounts of culture supernatants of HaCaT cells at 48 h were determined by ELISA. Data are expressed as means \pm SD ($n = 6$). ** $p < 0.01$, *** $p < 0.001$ significantly different from control. † $p < 0.05$, †† $p < 0.01$, ††† $p < 0.001$ significantly different from 5-FU alone.




Fig. 7. Effects of NAC on the levels of production of ROS and pro-inflammatory cytokines in 5-FU treated HaCaT cells. HaCaT cells was incubated with 5-FU in the presence or absence of NAC for 24 h (A–D) and 48 h (E–G). (A) ROS assay. Results are presented as the means \pm SD of three independent experiments. (B–G) IL-1 β , IL-6, and TNF- α levels were determined by real-time PCR. Bars represent mRNA expression normalized to GAPDH and relative to control. Data are expressed as means \pm SD ($n = 6$). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, significantly different from control. † $p < 0.01$, significantly different from 5-FU alone.

contributed to revise manuscript. All the authors read and approved the final manuscript.

Acknowledgments

This work was supported by Grants-in-Aid for Scientific Research (JSPS KAKENHI Grant Numbers JP18K09911 and JP21K10255) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Abbreviations

ARE	antioxidant response element
5-FU	5-fluorouracil
HaCaT cell	immortalized human keratinocyte cell line
HO-1	heme oxygenase 1

References

- | | |
|----------------|---|
| IL-1 β | interleukin (IL)-1 β |
| IL-6 | interleukin (IL)-6 |
| IL-10 | interleukin (IL)-10 |
| NAC | N-acetylcysteine |
| NAD | nicotinamide adenine dinucleotide |
| NF- κ B | nuclear factor kappa B |
| NQO-1 | NAD(P)H and quinone oxidoreductase 1 |
| Nrf2 | nuclear factor erythroid 2-related factor 2 |
| ROS | reactive oxygen species |
| SIRT-1 | silent information regulator transcript-1 |
| TNF- α | tumor necrosis factor- α |
- Conflict of Interest**
- No potential conflicts of interest were disclosed.
- Curra M, Soares Junior LAV, Martins MD, Santos PSDS. Chemotherapy protocols and incidence of oral mucositis. An integrative review. *Einstein (São Paulo)* 2018; **16**: eRW4007.
 - Rashid S, Ali N, Nafees S, Ahmad ST, Hasan SK, Sultana S. Abrogation of 5-fluorouracil induced renal toxicity by bee propolis via targeting oxidative stress and inflammation in Wistar rats. *J Pharm Pharmacol* 2013; **7**: 189–194.
 - Epstein JB, Robertson M, Emerton S, Phillips N, Stevenson-Moore P. Quality of life and oral function in patients treated with radiation therapy for head and neck cancer. *Head Neck* 2001; **23**: 389–398.
 - Sonis ST. A biological approach to mucositis. *J Support Oncol* 2004; **2**: 21–36.
 - Henke M, Alfonsi M, Foa P, et al. Palifermin decreases severe oral mucositis of patients undergoing postoperative radiochemotherapy for head and neck cancer: a randomized, placebo-controlled trial. *J Clin Oncol* 2011; **29**: 2815–2820.
 - Chang JW, Park KH, Hwang HS, Shin YS, Oh YT, Kim CH. Protective effects of Korean red ginseng against radiation-induced apoptosis in human HaCaT keratinocytes. *J Radiat Res* 2014; **55**: 245–256.
 - Sharma P, Jha AB, Dubey RS, Pessarakli M. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *J Bot* 2012; **2012**: 217037.
 - Rogers H, Munné-Bosch S. Production and scavenging of reactive oxygen species and redox signaling during leaf and flower senescence: similar but different. *Plant Physiol* 2016; **171**: 1560–1568.
 - Loboda A, Damulewicz M, Pyza E, Jozkowicz A, Dulak J. Role of Nrf2/HO-1 system in development, oxidative stress response and diseases: an evolutionarily conserved mechanism. *Cell Mol Life Sci* 2016; **73**: 3221–3247.
 - Yang HL, Lin SW, Lee CC, et al. Induction of Nrf2-mediated genes by *Antrodia salmonea* inhibits ROS generation and inflammatory effects in lipopolysaccharide-stimulated RAW264.7 macrophages. *Food Funct* 2015; **6**: 230–241.
 - Fu Q, Cui Q, Yang Y, et al. Effect of resveratrol dry suspension on immune function of piglets. *Evid Based Complement Alternat Med* 2018; **2018**: 5952707.
 - Orihuela-Campos RC, Tamaki N, Mukai R, et al. Biological impacts of resveratrol, quercetin, and N-acetylcysteine on oxidative stress in human gingival fibroblasts. *J Clin Biochem Nutr* 2015; **56**: 220–227.
 - Xiao Q, Zhu W, Feng W, et al. A review of resveratrol as a potent chemoprotective and synergistic agent in cancer chemotherapy. *Front Pharmacol* 2019; **9**: 1534.
 - Dos Santos Filho EX, da Silva ACG, de Ávila RI, et al. Chemopreventive effects of FITOPROT against 5-fluorouracil-induced toxicity in HaCaT cells. *Life Sci* 2018; **193**: 300–308.
 - Partridge M, Chantry D, Turner M, Feldmann M. Production of interleukin-1 and interleukin-6 by human keratinocytes and squamous cell carcinoma cell lines. *J Invest Dermatol* 1991; **96**: 771–776.
 - Konur A, Schulz U, Eissner G, Andreesen R, Holler E. Interferon (IFN)-gamma is a main mediator of keratinocyte (HaCaT) apoptosis and contributes to autocrine IFN-gamma and tumour necrosis factor-alpha production. *Br J*
 - Dermatol* 2005; **152**: 1134–1142.
 - Kim DR, Kim J, Oh JY, Kim HY, Kim YJ, Chang MS. Protective effect of *Salvia miltiorrhiza* Bunge on 5-fluorouracil-induced oral mucositis. *Int J Mol Med* 2017; **40**: 39–46.
 - Kang MA, So EY, Simons AL, Spitz DR, Ouchi T. DNA damage induces reactive oxygen species generation through the H2AX-Nox1/Rac1 pathway. *Cell Death Dis* 2012; **3**: e249.
 - Barbosa MM, de Araújo Júnior RF, et al. Telmisartan modulates the oral mucositis induced by 5-Fluorouracil in hamsters. *Front Physiol* 2018; **9**: 1204.
 - Sonis ST. Pathobiology of oral mucositis: novel insights and opportunities. *J Support Oncol* 2007; **5 (9 Suppl 4)**: 3–11.
 - Harris DJ. Cancer treatment-induced mucositis pain: strategies for assessment and management. *Ther Clin Risk Manag* 2006; **2**: 251–258.
 - Leonard SS, Xia C, Jiang BH, et al. Resveratrol scavenges reactive oxygen species and effects radical-induced cellular responses. *Biochem Biophys Res Commun* 2003; **309**: 1017–1026.
 - Kurutas EB. The importance of antioxidants which play the role in cellular response against oxidative/nitrosative stress: current state. *Nutr J* 2016; **15**: 71.
 - Singh N, Agrawal M, Doré S. Neuroprotective properties and mechanisms of resveratrol in *in vitro* and *in vivo* experimental cerebral stroke models. *ACS Chem Neurosci* 2013; **4**: 1151–1162.
 - Shen C, Cheng W, Yu P, et al. Resveratrol pretreatment attenuates injury and promotes proliferation of neural stem cells following oxygen-glucose deprivation/reoxygenation by upregulating the expression of Nrf2, HO-1 and NQO-1 *in vitro*. *Mol Med Rep* 2016; **14**: 3646–3654.
 - Dong W, Wang X, Bi S, et al. Inhibitory effects of resveratrol on foam cell formation are mediated through monocyte chemotactic protein-1 and lipid metabolism-related proteins. *Int J Mol Med* 2014; **33**: 1161–1168.
 - Dong WW, Liu YJ, Lv Z, et al. Lung endothelial barrier protection by resveratrol involves inhibition of HMGB1 release and HMGB1-induced mitochondrial oxidative damage via an Nrf2-dependent mechanism. *Free Radic Biol Med* 2015; **88 (Pt B)**: 404–416.
 - Chao SC, Chen YJ, Huang KH, et al. Induction of sirtuin-1 signaling by resveratrol induces human chondrosarcoma cell apoptosis and exhibits anti-tumor activity. *Sci Rep* 2017; **7**: 3180.
 - Ren Z, He H, Zuo Z, Xu Z, Wei Z, Deng J. The role of different SIRT1-mediated signaling pathways in toxic injury. *Cell Mol Biol Lett* 2019; **24**: 36.
 - Mukaigasa K, Nguyen LT, Li L, Nakajima H, Yamamoto M, Kobayashi M. Genetic evidence of an evolutionarily conserved role for Nrf2 in the protection against oxidative stress. *Mol Cell Biol* 2012; **32**: 4455–4461.
 - Nguyen T, Nioi P, Pickett CB. The Nrf2-antioxidant response element signaling pathway and its activation by oxidative stress. *J Biol Chem* 2009; **284**: 13291–13295.
 - Li L, Dong H, Song E, Xu X, Liu L, Song Y. Nrf2/ARE pathway activation, HO-1 and NQO-1 induction by polychlorinated biphenyl quinone is associated with reactive oxygen species and PI3K/AKT signaling. *Chem Biol Interact* 2014; **209**: 56–67.

- 33 Wang S, Wang J, Zhao A, Li J. SIRT1 activation inhibits hyperglycemia-induced apoptosis by reducing oxidative stress and mitochondrial dysfunction in human endothelial cells. *Mol Med Rep* 2017; **16**: 3331–3338.
- 34 Han MK, Song EK, Guo Y, Ou X, Mantel C, Broxmeyer HE. SIRT1 regulates apoptosis and nanog expression in mouse embryonic stem cells by controlling p53 subcellular localization. *Cell Stem Cell* 2008; **2**: 241–251.
- 35 Yeung F, Hoberg JE, Ramsey CS, et al. Modulation of NF-κB-dependent transcription and cell survival by the SIRT1 deacetylase. *EMBO J* 2004; **23**: 2369–2380.
- 36 Rada P, Pardo V, Mobasher MA, et al. SIRT1 controls acetaminophen hepatotoxicity by modulating inflammation and oxidative stress. *Antioxid Redox Signal* 2018; **28**: 1187–1208.
- 37 Chan SH, Hung CH, Shih JY, et al. SIRT1 inhibition causes oxidative stress and inflammation in patients with coronary artery disease. *Redox Biol* 2017; **13**: 301–309.
- 38 Huang KP, Huang J, Xie X, et al. Sirt1 resists advanced glycation end products-induced expressions of fibronectin and TGF-β1 by activating the Nrf2/ARE pathway in glomerular mesangial cells. *Free Radic Biol Med* 2013; **65**: 528–540.
- 39 Ding YW, Zhao GJ, Li XL, et al. SIRT1 exerts protective effects against paraquat-induced injury in mouse type II alveolar epithelial cells by deacetylating NRF2 *in vitro*. *Int J Mol Med* 2016; **37**: 1049–1058.
- 40 Wardyn JD, Ponsford AH, Sanderson CM. Dissecting molecular cross-talk between Nrf2 and NF-κB response pathways. *Biochem Soc Trans* 2015; **43**: 621–626.
- 41 Zhang Z, Liang Z, Li H, et al. Perfluorocarbon reduces cell damage from blast injury by inhibiting signal paths of NF-κB, MAPK and Bcl-2/Bax signaling pathway in A549 cells. *PLoS One* 2017; **12**: e0173884.
- 42 Albenis BC. What is nuclear factor kappa B (NF-κB) doing in and to the mitochondrion? *Front Cell Dev Biol* 2019; **7**: 154.
- 43 Pervaiz S. Resveratrol: from grapevines to mammalian biology. *FASEB J* 2003; **17**: 1975–1985.



This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).