

REVIEW

Varying clinical presentations of nontuberculous mycobacterial disease : Similar to but different from tuberculosis

Tsutomu Shinohara^{1,2,3}, Shun Morizumi^{1,2}, and Kenya Sumitomo²

¹Department of Community Medicine for Respiriology, Graduate School of Biomedical Sciences, Tokushima University, Tokushima, Japan, ²Division of Internal Medicine, Japan Agricultural Cooperatives Kochi Hospital, Kochi, Japan, ³Department of Clinical Investigation, National Hospital Organization Kochi Hospital, Kochi, Japan

Abstract : The incidence rate of pulmonary nontuberculous mycobacterial disease (PNTMD) in Japan is the highest among major industrialized nations. Although the typical clinical course and radiological manifestations of PNTMD are different from those of pulmonary tuberculosis (TB), confusion about these mycobacterial diseases leads to a diagnostic pitfall. Diagnostic challenges include the coexistence of *Mycobacterium tuberculosis* (MTB) and nontuberculous mycobacteria (NTM), false positives for NTM in MTB nucleic acid amplification tests, microbial substitution, and abnormal radiological manifestations caused by NTM. Features of extrapulmonary NTM diseases, such as pleurisy, vertebral osteomyelitis, and disseminated disease, are different from the corresponding tuberculous diseases. Moreover, the immunological background of the patient (status of human immunodeficiency virus infection with or without antiviral therapy, continuation or discontinuation of immunosuppressive therapy, use of immune checkpoint inhibitor, pregnancy and delivery, etc.) influences the pathophysiology of mycobacterial diseases. This review describes the varying clinical presentations of NTM disease with emphasis on the differences from TB. *J. Med. Invest.* 68 : 220-227, August, 2021

Keywords : nontuberculous mycobacteria, tuberculosis, acid-fast bacteria tests, extrapulmonary diseases, immune reconstitution inflammatory syndrome

INTRODUCTION

The incidence rate of pulmonary nontuberculous mycobacterial disease (PNTMD) in Japan during 2014 was estimated to be 14.7 cases per 100,000 person-years, exceeding that of culture-positive pulmonary tuberculosis (PTB) for the first time (1). This rate was the highest among major industrialized nations (2). Patients with PNTMD generally have a long disease course and the estimated PNTMD prevalence in Japan is greater than 110 cases per 100,000 persons, suggesting the increasing importance of cooperation between respiratory specialists and general physicians.

On the other hand, many physicians in Japan may lack knowledge about acid-fast bacteria (AFB) tests due to the decreasing number of tuberculosis (TB) patients. Furthermore, when physicians with little experience in treating TB patients provide medical care to patients with nontuberculous mycobacteria (NTM), they may confuse these mycobacterial diseases, leading to a diagnostic pitfall.

In addition, the general use of biological products for autoimmune and malignant diseases has led to an increase of mycobacterial diseases in patients whose immune response was altered by therapeutic agents. Therefore, physicians need to be reminded that they may encounter mycobacterial disease regardless of their specialty. This review describes the varying clinical presentations of NTM disease, and discusses the challenges of AFB tests and points to keep in mind when providing medical care to patients with NTM.

DIAGNOSTIC CHALLENGES IN PULMONARY MYCOBACTERIAL DISEASES

The typical clinical course and radiological manifestations of PNTMD are different from those of PTB, and the differential diagnosis between these infections can be simplified by identifying the mycobacterial species detected from the patient and use of supplementary methods such as *Mycobacterium tuberculosis* (MTB) antigen specific interferon- γ release assay (IGRA) and measurement of antibodies to the glycopeptidolipid (GPL) core antigen specific to *Mycobacterium avium* complex (MAC) (3-5). However, diagnostic pitfalls may cause physicians to make an incorrect initial diagnosis. Some of the diagnostic challenges of pulmonary mycobacterial diseases are as follows :

Erroneous diagnosis of multidrug-resistant tuberculosis (MDR-TB) caused by the presence of NTM

In Japan, MDR-TB, defined as the disease caused by MTB resistant to isoniazid and rifampicin, accounts for less than 1% of new TB cases (6, 7). As patients with MDR-TB are refractory to treatment, information about drug-susceptibility tests (DST) is important in clinical practice (6). However, the results of DST may not always reflect the subsequent clinical course. The diagnosis of MDR-TB was reported to be incorrect in 9 (13%) of 70 patients possibly due to laboratory-related errors (8). Situations that are considered to be laboratory-related errors include a good treatment course with standard regimens, disagreement with the DST results on resubmitted specimens, detection of MDR-TB from untreated inactive lesions, and a lack of tuberculous lesions. Coexistence of NTM with MTB in sputum, cross-contamination, and specimen mislabeling could be the cause (8, 9). Misdiagnosis of MDR-TB may cause serious harm to the patient because it will result in the unnecessary abandonment of an effective treatment. It is thus important to carefully observe the solid media to distinguish between colonies of NTM (smooth type) and MTB (rough type), and share information about the

Received for publication December 8, 2020 ; accepted March 22, 2021.

Address correspondence and reprint requests to Tsutomu Shinohara, MD, PhD., Department of Community Medicine for Respiriology, Graduate School of Biomedical Sciences, Tokushima University, 3-18-15 Kuramoto-cho, Tokushima 770-8503, Japan and Fax : +81-88-863-2186.

patient's background and clinical course between laboratory staff and the physicians to avoid misdiagnosis.

False positives for NTM in MTB nucleic acid amplification tests (NAATs)

An important part of the initial assessment of suspected pulmonary mycobacterial disease is the determination of whether it is PTB exhibiting human-to-human transmission. Currently, MTB NAATs are mainly used for rapid diagnosis (7 kits are currently available in Japan), and their sensitivities in patients with sputum smear-positive or negative PTB are 94-100% and 62-80%, respectively (10, 11). They also have excellent specificity (99-100%) (10, 11), but yield false positives for several NTM due to the homology of the gene sequence of the amplification site.

For example, TRCRapid M. TB (Tosoh Bioscience, Tokyo, Japan) utilizing a transcription-reverse transcription concerted (TRC) method that allows rapid detection of the target RNA yields false positives for several NTM, including *Mycobacterium shinjukuense* (12-21). If NAAT is positive in a suspected TB patient but the subsequent clinical course is inconsistent with TB, the isolate should be identified by the newly developed matrix-assisted laser desorption ionization-time of flight mass spectrometry or sequence analysis (22). False-positive reactions in MTB NAATs may occur or have already occurred for unidentified NTM species without being noticed.

Microbial substitution

Most of the MAC or *Mycobacterium abscessus* complex (MABC) isolates re-identified after negative sputum culture conversion obtained by anti-NTM chemotherapy are the result of reinfections with mycobacteria of different genotypes (23, 24). Furthermore, in the Japanese 2014 survey, the incidence rate of MABC disease increased five-fold in 7 years and microbial substitution of MAC to MABC is regarded as one of the reasons (1). MABC is classified into *M. abscessus* ssp. *abscessus*, *M. abscessus* ssp. *bolletii* and *M. abscessus* ssp. *massiliense*. The ribosomal methyl transferase gene *erm* (41) confers inducible macrolide resistance (susceptible on day 3 but resistant on day 14 of DST) in *M. abscessus* ssp. *abscessus* and *bolletii*. However *M. abscessus* ssp. *massiliense* is susceptible to clarithromycin even after prolonged incubation because its *erm* (41) does not work (24-27). In Japan, the rapidly growing mycobacteria identification kit using nucleic acid chromatography (KANEKA Corporation, Osaka, Japan) has made it possible to evaluate MABC subspecies and the *erm* (41) genotype in clinical practice. Management of NTM disease requires regular assessment of microbial substitution and reconsideration of treatment plans.

Abnormal radiological manifestations caused by NTM : Not what it seems

PNTMD generally presents as bronchiectasis with clustered centrilobular nodules (nodular bronchiectatic disease) or upper-lobe cavitory lesions (fibrocavitary disease). However, it occasionally develops as unexpected lesions.

1. Pneumonia-like shadows

The clinical manifestations of pulmonary mycobacterial diseases are affected by the immunological background of the host. For example, significant differences have been reported in chest radiographic and computed tomography (CT) findings of TB depending on the status of human immunodeficiency virus (HIV) infection and CD4 T cells (28, 29). TB can develop in the early stage of HIV infection, but NTM disease usually develops in the late stage (30, 31). Therefore, in HIV-positive patients with NTM disease, granulomas are not sufficiently formed and exudative lesions become predominant (32). On the other hand, rapidly progressive pulmonary consolidation due to granulomatous

inflammation in a patient with PNTMD without immunodeficiency was reported and similar shadows have been found in patients with rheumatoid arthritis (33-34). In such cases, an excessive immune response is assumed to be the principal pathological condition and steroid therapy is usually effective (33-34).

2. Miliary pulmonary opacities

A miliary pattern, randomly distributed nodular opacities, on chest radiographs is generally caused by a systemic disorder such as hematogenous spread of infection or cancer, and sarcoidosis. TB is a well-known infectious disease that can present with a miliary pattern (35), but NTM disease may also exhibit the same pattern. In an autopsy-based study, a miliary pattern was found in cases of disseminated NTM disease with pulmonary involvement, but not in cases of PNTMD alone (36).

However, miliary opacities are known to be observed in patients with hypersensitivity pneumonitis (HP), or extrinsic allergic alveolitis, and inhalation of aerosols containing NTM (mainly MAC) can induce a similar clinical presentation. Hot tub lung (HTL) is a typical disease of this condition and pathologically a granulomatous lung disease (37-39). Like HP, HTL shows apparent improvement either spontaneously or in response to glucocorticoid therapy after avoiding exposure to the pathogen without anti-NTM chemotherapy (39). The fact that bacillus Calmette-Guerin (BCG) immunotherapy for bladder cancer sometimes induces systemic granulomatous inflammation supports the hypothesis that HTL is an allergic reaction rather than an infection (40, 41). In addition, the possibility of endobronchial spread of NTM (*M. xenopi*) was reported in an immunocompromised patient (42).

3. Large pulmonary solitary mass

NTM, especially MAC, infections sometimes cause solitary pulmonary nodules (SPN) (43). In Japan, the proportion of *M. intracellulare* cases is high in the southwestern region (44), but SPN are mainly due to *M. avium* infections. SPN in patients with NTM infection are less likely to cause satellite lesions and calcification than those in patients with MTB infection. Therefore, this type of disease may not be diagnosed until surgical resection is performed for suspected lung cancer (43). In addition, granulomas and expanded bronchi with purulent exudate can aggregate together to form a large mass. We previously reported a lobulated mass with marked 18-fluorodeoxyglucose (FDG) uptake, mimicking pulmonary mucinous cystadenocarcinoma due to *Mycobacterium kansasii* infection (45). As the presentations of PNTMD are protean, NTM infection should not be excluded from the differential diagnosis of pulmonary disease simply based on radiological manifestations alone.

FEATURES OF EXTRAPULMONARY NTM DISEASES THAT ARE DIFFERENT FROM THE CORRESPONDING TUBERCULOUS DISEASES

As MTB survives only in humans and exhibits aerial transmission, it is not directly transmitted into the extrapulmonary organs from outside of the body. During primary infection, hematogenous and lymphatic spread of MTB to extrapulmonary organs occurs before adaptive T-cell immunity is firmly established. There are two presentations: early-onset following primary infection and endogenous recurrence from long-surviving MTB in microgranulomatous lesions (46). However, NTM residues in natural environments (soil and water) and can transmit through the skin, gastrointestinal tract, and sharp injuries to form extrapulmonary lesions.

Pleurisy

Tuberculous pleurisy that follows primary infection (primary TB pleurisy) is the most common extrapulmonary TB in Japan, and its pathogenesis is considered to be a delayed-type allergic reaction to TB antigens in the pleural cavity (47). Multiple cytokines released mainly from immune cells increase the permeability of the pleural capillaries and cause pleural effusion (48, 49). This condition can resolve spontaneously, but recurrence was observed in approximately half of untreated cases resulting in serious presentation (50, 51). However, in asymptomatic patients with pleural thickening suspected to be caused by primary TB pleurisy, antituberculous chemotherapy is usually not administered. We previously reported the usefulness of FDG positron emission tomography/CT evaluation in such a case (52).

On the other hand, primary NTM pleurisy (i.e., without distinct pulmonary involvement) is rare, probably because NTM are environmental mycobacteria inducing desensitization in hosts before its infection (53). NTM pleurisy may be caused by a direct extension of infection from pulmonary lesions into the pleural cavity and is often intractable (54-56). Endoscopic examination to diagnose PNTMD was reported to cause NTM pleuritis with pneumothorax probably due to the rupture of a subpleural lesion during the procedure (20, 57). Recently, Furuchi *et al.* reported a case of PNTMD with pleural effusion, suggesting that evaluation of pleural fluid anti-GPL-core IgA antibody can be a supplementary diagnostic method for the pleural involvement of MAC (58).

Vertebral osteomyelitis

Vertebral osteomyelitis (VO) is now an uncommon clinical presentation of TB, but tuberculous VO was reported to be increasing in the elderly. Patients with bone and joint TB usually have few typical symptoms and inflammatory markers in patients are often not elevated. In addition, age-related changes in the vertebra, such as spinal joint deformities and osteoporosis, can affect radiological findings, leading to oversights or delays in the diagnosis of tuberculous VO (59).

NTM also cause VO, with about half of the cases occurring in otherwise healthy patients (60-64). Blunt-force trauma was reported to be a risk factor for VO due to mycobacterial infection, and it is hypothesized that macrophages which have phagocytosed NTM migrate to the injury site for tissue restoration but release the pathogen resulting in the development of VO (62). An interesting case of *M. abscessus* ssp. *abscessus* disease was recently reported, which was initially diagnosed as primary pleurisy, but later turned out to be disease progression from VO (65). The major pathogen of VO, including cases with exudative pleural effusion, is *Staphylococcus aureus* (66, 67), but NTM infection should be considered, especially in patients with a history of pre-disposing trauma.

Disseminated mycobacterial disease

Disseminated TB is defined as the hematogenous spread of tuberculous infection involving multiple noncontiguous sites (35). Miliary TB is a term that refers to miliary granulomas under the same conditions generally with innumerable tiny spots on a chest radiograph (68).

Disseminated NTM disease has been observed in adults with advanced HIV infection and in cases of severe cellular immune dysfunction caused by immunosuppressive therapy such as after hematopoietic stem cell and organ transplantation (30, 31, 69). In recent years, neutralizing anti-interferon- γ autoantibody has received a great deal of attention as a new etiology of this condition (70-72). It should be noted that the main primary site of transmission of disseminated MAC (mainly *M. avium*) disease

is the intestinal mucosa and this condition does not necessarily have a preceding lung lesion (73). Blood cultures play an important role in the diagnosis (74) and rapidly growing mycobacteria may be detected in commonly used aerobic culture bottles. However, BACTEC Myco/F Lytic culture vials are more suitable for the detection of mycobacterium, especially for slowly growing species, including MAC (75). NTM can also be the causative pathogens of catheter-related bloodstream infections (76).

MYCOBACTERIAL DISEASES ASSOCIATED WITH IMMUNE RECONSTITUTION INFLAMMATORY SYNDROME (IRIS)

IRIS is a condition noted in some cases of rapid reconstruction or activation of immune function, which increases the immunoinflammatory response to antigens mainly derived from microorganisms. It typically causes the development of a new opportunistic infection (unmasking IRIS) or paradoxical clinical exacerbation of a pre-existing infection (paradoxical IRIS) (77). Mycobacterial diseases in representative clinical settings of IRIS are described below.

Introduction of antiretroviral therapy (ART)

Paradoxical exacerbation of existing TB during antituberculous chemotherapy was observed in 36% of HIV-positive patients who were additionally treated using ART. The median duration of antituberculous chemotherapy before the onset of paradoxical TB-IRIS was 8 weeks (interquartile range 5.5-11.5 weeks), whereas the median duration of ART was only 4 weeks (interquartile range 2-10 weeks). In addition to the exacerbation of the preceding lesions, new lesions may develop in extrapulmonary organs, including the central nervous system (78, 79). In this setting, NTM disease is often recognized as unmasking IRIS and superficial lymphadenitis is common. Unlike disseminated NTM disease, lesions may be localized and the positive rate on blood culture is low (79).

Discontinuation of TNF inhibitors and immunosuppressive drugs

Tumor necrosis factor (TNF)- α is required for the formation and maintenance of granuloma, which confine microbial pathogens difficult to eradicate. Therefore, administration of TNF inhibitors to patients with autoimmune diseases, such as rheumatoid arthritis, can induce granulomatous infections, including activation of latent TB (80, 81). Discontinuation of anti-TNF therapy at the onset of TB can cause paradoxical exacerbation (82). According to the Japanese guidelines, if the susceptibility of MTB to anti-TB drugs is known and TNF inhibitors can be safely administered, continuation of the treatment using a biological product during antituberculous chemotherapy should be considered (83).

Regarding NTM diseases associated with anti-TNF therapy, the majority of patients were administered either corticosteroids or methotrexate at the same time, and extrapulmonary lesions were frequently observed (84). The paradoxical exacerbation of NTM diseases in patients with autoimmune diseases while undergoing dose reduction, or after the discontinuation of TNF inhibitors or immunosuppressive drugs has also been reported (85-87). In general, NTM disease is refractory to treatment, making it difficult to notice the paradoxical exacerbation. However, in the above cases, persistent infection with NTM was excluded by AFB tests. Moreover, the introduction of steroid therapy or increase in the dose of glucocorticoid and immunosuppressive drug was effective.

Introduction of immune checkpoint inhibitors (ICIs)

The development of a new opportunistic infection during ICI therapy has become a concern and many TB cases were considered to be unmasking IRIS. Therefore, confirmation of a negative IGRA result prior to ICI administration may be beneficial, similar to before the administration of TNF inhibitors (88). Approximately half of the cases of unmasking TB during ICI therapy were detected before the onset of subjective symptoms by CT performed for the purpose of evaluation of ICI therapy against malignancy (89). The paradoxical exacerbation of existing TB during nivolumab therapy was also reported in a lung cancer patient (90). Regarding NTM disease during ICI therapy, there are only a few reported cases from Japan. Nodular bronchiectatic disease developed in advanced lung cancer patients previously treated using cytotoxic chemotherapy and thoracic radiotherapy (91). Conversely, there was a case in which the existing pulmonary *M. abscessus* ssp. *massiliense* disease was improved by ICI therapy for advanced lung cancer (92). The accumulation of cases of NTM disease in this setting is necessary in order to improve the management of ICI-induced immune-mediated events.

Recovery from immunosuppression by treatment for the underlying disease

Paradoxical clinical exacerbation of TB has been recognized, usually after initial improvement, in otherwise healthy patients receiving antituberculous chemotherapy, and is referred to as a paradoxical response or paradoxical reaction (PR) (93). This phenomenon may be attributed to an excessive immune response to massive antigen release from dying MTB coincided with recovery from TB-induced immunosuppression. In general, the duration of antituberculous chemotherapy before paradoxical exacerbation is longer in HIV-negative patients than in HIV-positive patients on ART (94).

Several cases of PR in otherwise healthy patients treated for NTM have also been reported, among which exacerbation of existing lesions and the development of new lesions were noted after negative conversion of the sputum AFB smear results without redetection of NTM (95, 96). These lesions gradually improved without changing the chemotherapy regimen.

Recovery from an immunosuppressive condition during pregnancy

Pregnancy is regarded as an immunosuppressive condition to maintain fetal tolerance and the Th1 response recovers rapidly from the suppressed state during the postpartum period resulting in the development of new infections including TB (unmasking IRIS) (97, 98). In postpartum TB cases, the frequencies of extrapulmonary and central nervous system involvement were reported to be approximately 90 and 70%, respectively. The mortality rate and frequency of sequelae were 38 and 14%, respectively (99). Postpartum tuberculosis may progress to PR without initial improvement on the same IRIS spectrum (100). To the best of our knowledge, there are no reports of postpartum PNTMD. However, many cases of postcesarean section wound infection caused by MABC have been reported, although the involvement of immune reconstitution has not been discussed (101, 102).

CONCLUSIONS

In the diagnosis and treatment of pulmonary mycobacterial disease, it is important to confirm that there is no discrepancy between the clinical course and the results of AFB tests. If necessary, new and/or existing samples must be examined to confirm the mycobacterial species. As the presentations of PNTMD

are protean, NTM infection should not be excluded from the differential diagnosis of pulmonary disease simply based on radiological manifestations alone. The development of extrapulmonary NTM diseases is different from that of corresponding tuberculous diseases because NTM are environmental mycobacteria that can transmit through the skin, gastrointestinal tract and sharps injuries to form extrapulmonary lesions. In addition, noninvasive trauma may be a risk factor for VO due to NTM infection. Moreover, it is necessary to pay sufficient attention to the immune function and its modifiers. As an opportunistic infection, NTM disease usually develops with more advanced immunosuppression than TB and is more likely to be disseminated. On the other hand, NTM disease associated with IRIS is less common and unmasking disease is often localized compared with TB. The paradoxical exacerbation of NTM disease as IRIS is usually responsive to the increase in immunosuppression.

REFERENCES

- Namkoong H, Kurashima A, Morimoto K, Hoshino Y, Hasegawa N, Ato M, Mitarai S : Epidemiology of pulmonary nontuberculous mycobacterial disease, Japan. *Emerg Infect Dis* 22 : 1116-1117, 2016
- Prevots DR, Marras TK : Epidemiology of human pulmonary infection with nontuberculous mycobacteria : a review. *Clin Chest Med* 36 : 13-34, 2015
- Mori T : Usefulness of interferon-gamma release assays for diagnosing TB infection and problems with these assays. *J Infect Chemother* 15 : 143-155, 2009
- Shibata Y, Horita N, Yamamoto M, Tsukahara T, Nagakura H, Tashiro K, Watanabe H, Nagai K, Nakashima K, Ushio R, Ikeda M, Narita A, Kanai A, Sato T, Kaneko T : Diagnostic test accuracy of anti-glycopeptidolipid-core IgA antibodies for *Mycobacterium avium* complex pulmonary disease : systematic review and meta-analysis. *Sci Rep* 6 : 29325, 2016, doi : 10.1038/srep29325
- Kitada S, Maekura R, Yoshimura K, Miki K, Miki M, Oshitani Y, Nishida K, Sawa N, Mori M, Kobayashi K : Levels of Antibody against Glycopeptidolipid Core as a Marker for Monitoring Treatment Response in *Mycobacterium avium* Complex Pulmonary Disease : a Prospective Cohort Study. *J Clin Microbiol* 55 : 884-892, 2017
- Dheda K, Gumbo T, Gandhi NR, Murray M, Theron G, Udwadia Z, Migliori GB, Warren R : Global control of tuberculosis : from extensively drug-resistant to untreatable tuberculosis. *Lancet Respir Med* 2 : 321-338, 2014
- Tuberculosis Research Committee (RYOKEN), Tokyo, Japan : Nationwide survey of anti-tuberculosis drug resistance in Japan. *Int J Tuberc Lung Dis* 19 : 157-162, 2015
- Nitta AT, Davidson PT, de Koning ML, Kilman RJ : Misdiagnosis of multidrug-resistant tuberculosis possibly due to laboratory-related errors. *JAMA* 276 : 1980-1983, 1996
- Ohji R, Shinohara T, Ohji H, Hatakeyama N, Iwamura S, Ogushi F : Pulmonary tuberculosis misdiagnosed as extensively drug-resistant tuberculosis due to selection of a nontuberculous mycobacterial colony from a preculture dish used for a drug-susceptibility test. *Respir Investig* 58 : 216-219, 2020
- Clarridge JE 3rd, Shawar RM, Shinnick TM, Plikaytis BB : Large-scale use of polymerase chain reaction for detection of *Mycobacterium tuberculosis* in a routine mycobacteriology laboratory. *J Clin Microbiol* 31 : 2049-2056, 1993
- Yoshida H, Onohara K, Tazawa T, Tsurinaga Y, Kurokawa M, Han Y, Tamura Y, Nagai T, Hashimoto S, Kawase I : Comparative study of the efficacy of the COBAS TaqMan

- and LAMP assay for the rapid diagnosis of tuberculosis. *Kekkaku* 88 : 727-733, 2013 (in Japanese)
12. Saito H, Iwamoto T, Ohkusu K, Otsuka Y, Akiyama Y, Sato S, Taguchi O, Sueyasu Y, Kawabe Y, Fujimoto H, Ezaki T, Butler R : *Mycobacterium shinjukuense* sp. nov., a slowly growing, non-chromogenic species isolated from human clinical specimens. *Int J Syst Evol Microbiol* 61(Pt 8) : 1927-1932, 2011
 13. Aono A, Kazumi Y, Maeda S, Azuma Y, Tsuchiya S, Iwamoto T, Nakanaga K, Hayakawa K, Saito H : Non-tuberculous mycobacterium strains that show positive test for identification kits of *M. tuberculosis* complex. *Kekkaku* 85 : 461-464, 2010 (in Japanese)
 14. Watanabe K, Shinkai M, Yamaguchi N, Shinoda M, Hara Y, Ishigatsubo Y, Kaneko T : *Mycobacterium shinjukuense* lung disease that was successfully treated with antituberculous drugs. *Intern Med* 52 : 2653-2655, 2013
 15. Takeda K, Ohshima N, Nagai H, Sato R, Ando T, Kusaka K, Kawashima M, Masuda K, Matsui H, Aono A, Chikamatsu K, Mitarai S, Ohta K : Six Cases of Pulmonary *Mycobacterium shinjukuense* infection at a single hospital. *Intern Med* 55 : 787-791, 2016
 16. Ishiguro T, Saitoh J, Horie R, Hayashi T, Ishizuka T, Tsuchiya S, Yasukawa K, Kido T, Nakaguchi Y, Nishibuchi M, Ueda K : Intercalation activating fluorescence DNA probe and its application to homogeneous quantification of a target sequence by isothermal sequence amplification in a closed vessel. *Anal Biochem* 314 : 77-86, 2003
 17. Takeda H, Hirose H, Ogura A, Kato Y, Sugiyama Y, Harada H, Fukuno Y, Yamamura T, Miyajima E : Comparison of Tosoh TRCRapid M.TB assay by transcription-reverse transcription concerted reaction (TRC) with Roche COBAS AMPLICOR assay by PCR and with culture for detection of *Mycobacterium tuberculosis* complex. *Rinsho Byori* 56 : 277-282, 2008 (in Japanese)
 18. Oshima K, Yokouchi H, Minemura H, Saito J, Tanino Y, Munakata M. Pulmonary infection caused by *Mycobacterium shinjukuense*. *Ann Am Thorac Soc* 12 : 958-959, 2015
 19. Hayashi M, Matsukura S, Funaki T, Inoue D, Kazumi Y, Mitarai S, Kokubu F, Shikama Y : Clarithromycin-resistant *Mycobacterium Shinjukuense* lung disease : case report and literature review. *Showa Univ J Med Sci* 28 : 373-377, 2016
 20. Taoka T, Shinohara T, Hatakeyama N, Iwamura S, Murase Y, Mitarai S, Ogushi F : *Mycobacterium Shinjukuense* pulmonary disease progressed to pleuritis after iatrogenic pneumothorax : a case report. *J Clin Tuberc Other Mycobact Dis* 19 : 100160, 2020, doi : 10.1016/j.jctube.2020.100160. eCollection
 21. Moon SM, Kim SY, Chung MJ, Lee SH, Shin SJ, Koh WJ : Nontuberculous mycobacterial lung disease caused by *Mycobacterium shinjukuense* : the first reported case in Korea. *Tuberc Respir Dis (Seoul)* 78 : 416-418, 2015
 22. Luo L, Cao W, Chen W, Zhang R, Jing L, Chen H, Yu F, Yue J : Evaluation of the VITEK MS knowledge base version 3.0 for the identification of clinically relevant *Mycobacterium* species. *Emerg Microbes Infect* 7 : 114, 2018, doi : 10.1038/s41426-018-0120-3
 23. Wallace RJ Jr, Brown-Elliott BA, McNulty S, Philley JV, Killingley J, Wilson RW, York DS, Shepherd S, Griffith DE : Macrolide/Azalide therapy for nodular/bronchiectatic *mycobacterium avium* complex lung disease. *Chest* 146 : 276-282, 2014
 24. Koh WJ, Jeong BH, Kim SY, Jeon K, Park KU, Jhun BW, Lee H, Park HY, Kim DH, Huh HJ, Ki CS, Lee NY, Kim HK, Choi YS, Kim J, Lee SH, Kim CK, Shin SJ, Daley CL, Kim H, Kwon OJ : Mycobacterial characteristics and treatment outcomes in *Mycobacterium abscessus* lung disease. *Clin Infect Dis* 64 : 309-316, 2017
 25. Nash KA, Brown-Elliott BA, Wallace RJ Jr : A novel gene, *erm(41)*, confers inducible macrolide resistance to clinical isolates of *Mycobacterium abscessus*, but is absent from *Mycobacterium chelonae*. *Antimicrob Agents Chemother* 53 : 1367-1376, 2009
 26. Choi GE, Shin SJ, Won CJ, Min KN, Oh T, Hahn MY, Lee K, Lee SH, Daley CL, Kim S, Jeong BH, Jeon K, Koh WJ : Macrolide treatment for *Mycobacterium abscessus* and *Mycobacterium massiliense* infection and inducible resistance. *Am J Respir Crit Care Med* 186 : 917-925, 2012
 27. Bastian S, Veziris N, Roux AL, Brossier F, Gaillard JL, Jarlier V, Cambau E : Assessment of clarithromycin susceptibility in strains belonging to the *Mycobacterium abscessus* group by *erm(41)* and *rrl* sequencing. *Antimicrob Agents Chemother* 55 : 775-781, 2011
 28. Haramati LB, Jenny-Avital ER, Alterman DD : Effect of HIV status on chest radiographic and CT findings in patients with tuberculosis. *Clin Radiol* 52 : 31-35, 1997
 29. Laissy JP, Cadi M, Boudiaf ZE, Casalino E, Crestani B, Bouvet E, Aubier M, Schouman-Claeys E : Pulmonary tuberculosis : computed tomography and high-resolution computed tomography patterns in patients who are either HIV-negative or HIV-seropositive. *J Thorac Imaging* 13 : 58-64, 1998
 30. Kirk O, Gatell JM, Mocroft A, Pedersen C, Proenca R, Brettle RP, Barton SE, Sudre P, Phillips AN, Lundgren JD for the EuroSIDA study group : Infections with *Mycobacterium tuberculosis* and *Mycobacterium avium* among HIV-infected patients after the introduction of highly active antiretroviral therapy. *Am J Respir Crit Care Med* 162(3 Pt 1) : 865-872, 2000
 31. The Center for Disease Control and Prevention. the National Institutes of Health, and the HIV Medicine Association of the Infectious Diseases Society of America : Guidelines for the Prevention and Treatment of Opportunistic Infections in Adults and Adolescents with HIV. (Downloaded from <https://aidsinfo.nih.gov/guidelines> on 11/01/2020)
 32. Fujita J : Various Clinical and Pathological Findings of Pulmonary Non-tuberculous Mycobacteriosis-from the Viewpoint of Granuloma Formation-. *JJSOG* 39 : 11-17, 2019(in Japanese)
 33. Waller EA, Roy A, Brumble L, Khor A, Johnson MM, Garland JL : The expanding spectrum of *Mycobacterium avium* complex-associated pulmonary disease : *Chest* 130 : 1234-1241, 2006
 34. Yuki M, Tokuda H, Nagai H, Moda M, Emoto N, Kasai S, Ookouchi Y : Three cases of rheumatoid arthritis complicated with nontuberculous mycobacterial disease showing extensive consolidation in chest imaging. *Open J Respir* 4 : e00096, 2020 (in Japanese)
 35. Khan FY : Review of literature on disseminated tuberculosis with emphasis on the focused diagnostic workup. *J Family Community Med* 26 : 83-91, 2019
 36. O'Connell ML, Birkenkamp KE, Kleiner DE, Folio LR, Holland SM, Olivier KN : Lung manifestations in an autopsy-based series of pulmonary or disseminated nontuberculous mycobacterial disease. *Chest* 141 : 1203-1209, 2012
 37. Fahim A, Khan MN : An unusual cause of miliary radiographic pattern. *BMJ Case Rep* 2011, doi : 10.1136/bcr.10.2.011.5060
 38. Zota V, Angelis SM, Fraire AE, McNamee C, Kielbasa S, Libraty DH : Lessons from *Mycobacterium avium* complex-associated pneumonitis : a case report. *J Med Case Rep* 2 : 152, 2008

39. Hanak V, Kalra S, Aksamit TR, Hartman TE, Tazelaar HD, Ryu JH : Hot tub lung : presenting features and clinical course of 21 patients. *Respir Med* 100 : 610-615, 2006
40. Israel-Biet D, Venet A, Sandron D, Ziza JM, Chretien J : Pulmonary complications of intravesical Bacille Calmette-Guérin immunotherapy. *Am Rev Respir Dis* 135 : 763-765, 1987
41. Tobiume M, Shinohara T, Kuno T, Mukai S, Naruse K, Hatakeyama N, Ogushi F : BCG-induced pneumonitis with lymphocytic pleurisy in the absence of elevated KL-6. *BMC Pulm Med* 14 : 35, 2014, doi : 10.1186/1471-2466-14-35.
42. Okano Y, Shinohara T, Imanishi S, Takahashi N, Naito N, Taoka T, Kadota N, Ogushi F : Miliary pulmonary nodules due to *Mycobacterium xenopi* in a steroid-induced immunocompromised patient successfully treated with chemotherapy : a case report. *BMC Pulm Med* 16 : 92, 2016, doi : 10.1186/s12890-016-0252-y.
43. Ose N, Maeda H, Takeuchi Y, Susaki Y, Kobori Y, Taniguchi S, Maekura R : Solitary pulmonary nodules due to nontuberculous mycobacteriosis among 28 resected cases. *Int J Tuberc Lung Dis* 20 : 1125-1129, 2016
44. Morimoto K, Hasegawa N, Izumi K, Among H, Uchimura K, Yoshiyama T, Hoshino Y, Kurashima A, Sokunaga J, Shibuya S, Shimojima M, Ato M, Mitarai S : A Laboratory-based Analysis of Nontuberculous Mycobacterial Lung Disease in Japan from 2012 to 2013. *Ann Am Thorac Soc* 14 : 49-56, 2017
45. Kadota N, Shinohara T, Naruse K, Hino H, Ogushi F : A lobulated mass in the left lower lobe : not what it seems. *Thorax* 74 : 519-520, 2019
46. Rodriguez-Takeuchi SY, Renjifo ME, Medina FJ : Extrapulmonary tuberculosis : pathophysiology and imaging findings. *Radiographics* 39 : 2023-2037, 2019
47. Barnes PF, Mistry SD, Cooper CL, Pirmez C, Rea TH, Modlin RL : Compartmentalization of a CD4+ T lymphocyte subpopulation in tuberculous pleuritis. *J Immunol* 142 : 1114-1119, 1989
48. Momi H, Matsuyama W, Inoue K, Kawabata M, Arimura K, Fukunaga H, Osame M : Vascular endothelial growth factor and proinflammatory cytokines in pleural effusions. *Respir Med* 96 : 817-822, 2002
49. Shaw JA, Diacon AH, Koegelenberg CFN : Tuberculous pleural effusion. *Respirology* 24 : 962-971, 2019
50. Patiala J : Initial tuberculous pleuritis in the Finnish armed forces in 1939-1945 with special reference to eventual post-pleuritic tuberculosis. *Acta Tuberc Scand* 36 (Suppl) : 1-57, 1954
51. Roper WH, Waring JJ : Primary serofibrinous pleural effusion in military personnel. *Am Rev Tuberc* 71 : 616-634, 1955
52. Shinohara T, Shiota N, Kume M, Hamada N, Naruse K, Ogushi F : Asymptomatic primary tuberculous pleurisy with intense 18-fluorodeoxyglucose uptake mimicking malignant mesothelioma. *BMC Infect Dis* 13 : 12, 2013, doi : 10.1186/1471-2334-13-12.
53. Kamiya H, Toyota E, Kobayashi N, Kudo K : A case of pulmonary *Mycobacterium kansasii* infection complicated with pleural effusion. *Kekkaku* 79 : 397-400, 2004 (in Japanese).
54. Ichiki H, Ueda S, Watanabe A, Sato C, Abe M : Nontuberculous pulmonary mycobacteriosis complicated by pleuritis. *Nihon Kokyuki Gakkai Zasshi* 49 : 885-889, 2011 (in Japanese)
55. Sado T, Nakamura Y, Kita H : Clinical analysis of nontuberculous mycobacterial infection complicated by pleurisy. *Kekkaku* 89 : 821-824, 2014 (in Japanese)
56. Ando T, Kawashima M, Matsui H, Takeda K, Sato R, Ohshima N, Nagai H, Kitani M, Hebisawa A, Ohta K : Clinical Features and Prognosis of Nontuberculous Mycobacterial Pleuritis. *Respiration* 96 : 507-513, 2018
57. Nicholson TT, Mutlu GM : Pneumothorax following bronchoalveolar lavage for the diagnosis of non-tuberculous mycobacterial infection - An "atypical" complication of bronchoscopy? *Arch Bronconeumol* 52 : 278-279, 2016
58. Furuuchi K, Morimoto K, Fujiwara K, Tanaka Y, Takemura T, Kurashima A, Yoshimori K, Ohta K, Sasaki Y : Mycobacterium avium Complex Pleuritis with Elevated Anti-glycopeptidolipid-core IgA Antibody Levels in Pleural Effusion. *Intern Med* 58 : 2577-2579, 2019
59. Maeda Y, Izawa K, Nabeshima T, Yonenobu K : Tuberculous spondylitis in elderly Japanese patients. *J Orthop Sci* 13 : 16-20, 2008
60. Kim CJ, Kim UJ, Kim HB, Park SW, Oh MD, Park KH, Kim NJ : Vertebral osteomyelitis caused by non-tuberculous mycobacteria : Predisposing conditions and clinical characteristics of six cases and a review of 63 cases in the literature. *Infect Dis (Lond)* 48 : 509-516, 2016
61. Sarria JC, Chutkan NB, Figueroa JE, Hull A : Atypical mycobacterial vertebral osteomyelitis : case report and review. *Clin Infect Dis* 26 : 503-505, 1998
62. Chan ED, Kong PM, Fennelly K, Dwyer AP, Iseman MD : Vertebral osteomyelitis due to infection with nontuberculous mycobacterium species after blunt trauma to the back : 3 examples of the principle of locus minoris resistentiae. *Clin Infect Dis* 32 : 1506-1510, 2001
63. Garcia DC, Sandoval-Sus J, Razzaq K, Young L : Vertebral osteomyelitis caused by *Mycobacterium abscessus*. *BMJ Case Rep* 2013, doi : 10.1136/bcr-2013-009597
64. Edwards C, Diveronica M, Abel E : Epidural abscess caused by *Mycobacterium abscessus*. *Am J Case Rep* 13 : 180-182, 2012
65. Kadota N, Shinohara T, Hino H, Goda Y, Murase Y, Mitarai S, Ogushi F : *Mycobacterium abscessus* ssp. *abscessus* infection progressing to empyema from vertebral osteomyelitis in an immunocompetent patient without pulmonary disease : a case report. *BMC Pulm Med* 19 : 100, 2019, doi : 10.1186/s12890-019-0860-4
66. Inoue S, Moriyama T, Horinouchi Y, Tachibana T, Okada F, Maruo K, Yoshiya S : Comparison of clinical features and outcomes of staphylococcus aureus vertebral osteomyelitis caused by methicillin-resistant and methicillin-sensitive strains. *Springerplus* 2 : 283, 2013, doi : 10.1186/2193-1801-2-283
67. Bass SN, Ailani RK, Shekar R, Gerblisch AA : Pyogenic vertebral osteomyelitis presenting as exudative pleural effusion : a series of five cases. *Chest* 114 : 642-647, 1998
68. Matsushima T : Miliary tuberculosis or disseminated tuberculosis. *Intern Med* 44 : 687, 2005
69. Henkle E, Winthrop KL : Nontuberculous mycobacteria infections in immunosuppressed hosts. *Clin Chest Med* 36 : 91-99, 2015
70. Wu UI, Holland SM : Host susceptibility to non-tuberculous mycobacterial infections. *Lancet Infect Dis* 15 : 968-980, 2015
71. Browne SK, Burbelo PD, Chetchotissakd P, Suputtamongkol Y, Kiertiburanakul S, Shaw PA, Kirk JL, Jutivorakool K, Zaman R, Ding L, Hsu AP, Patel SY, Olivier KN, Lulitanond V, Mootsikapun P, Anunnatsiri S, Angkasekwinai N, Sathapatayavongs B, Hsueh PR, Shieh CC, Brown MR, Thongnoppakhun W, Claypool R, Sampaio EP, Thepthai C, Waywa D, Dacombe C, Reizes Y, Zelazny AM, Saleeb P, Rosen LB, Mo A, Iadarola M, Holland SM : Adult-onset immunodeficiency in Thailand and Taiwan. *N Engl J Med*

- 367 : 725-734, 2012
72. Aoki A, Sakagami T, Yoshizawa K, Shima K, Toyama M, Tanabe Y, Moro H, Aoki N, Watanabe S, Koya T, Hasegawa T, Morimoto K, Kurashima A, Hoshino Y, Trapnell BC, Kikuchi T : Clinical Significance of Interferon- γ Neutralizing Autoantibodies Against Disseminated Nontuberculous Mycobacterial Disease. *Clin Infect Dis* 66 : 1239-1245, 2018
 73. Horsburgh CR Jr : The pathophysiology of disseminated *Mycobacterium avium* complex disease in AIDS. *J Infect Dis* 179 (Suppl 3) : S461-465, 1999
 74. Yagupsky P, Menegus MA : Cumulative positivity rates of multiple blood cultures for *Mycobacterium avium*-intracellulare and *Cryptococcus neoformans* in patients with the acquired immunodeficiency syndrome. *Arch Pathol Lab Med* 114 : 923-925, 1990
 75. Vetter E, Torgerson C, Feuker A, Hughes J, Harmsen S, Schleck C, Horstmeier C, Roberts G, Cockerill F 3rd : Comparison of the BACTEC MYCO/F Lytic bottle to the isolator tube, BACTEC Plus Aerobic F/bottle, and BACTEC Anaerobic Lytic/10 bottle and comparison of the BACTEC Plus Aerobic F/bottle to the Isolator tube for recovery of bacteria, mycobacteria, and fungi from blood. *J Clin Microbiol* 39 : 4380-4386, 2001
 76. Tagashira Y, Kozai Y, Yamasa H, Sakurada M, Kashiya T, Honda H : A cluster of central line-associated bloodstream infections due to rapidly growing nontuberculous mycobacteria in patients with hematologic disorders at a Japanese tertiary care center : an outbreak investigation and review of the literature. *Infect Control Hosp Epidemiol* 36 : 76-80, 2015
 77. Haddow LJ, Easterbrook PJ, Mosam A, Khanyile NG, Parboosing R, Moodley P, Moosa MY : Defining immune reconstitution inflammatory syndrome : evaluation of expert opinion versus 2 case definitions in a South African cohort. *Clin Infect Dis* 49 : 1424-1432, 2009
 78. Narita M, Ashkin D, Hollender ES, Pitchenik AE : Paradoxical worsening of tuberculosis following antiretroviral therapy in patients with AIDS. *Am J Respir Crit Care Med* 158 : 157-161, 1998
 79. Lawn SD, Bekker LG, Miller RF : Immune reconstitution disease associated with mycobacterial infections in HIV-infected individuals receiving antiretrovirals. *Lancet Infect Dis* 5 : 361-373, 2005
 80. Wallis RS, Broder MS, Wong JY, Hanson ME, Beenhouwer DO : Granulomatous infectious diseases associated with tumor necrosis factor antagonists. *Clin Infect Dis* 38 : 1261-1265, 2004
 81. Winthrop KL : Risk and prevention of tuberculosis and other serious opportunistic infections associated with the inhibition of tumor necrosis factor. *Nat Clin Pract Rheumatol* 2 : 602-10, 2006
 82. Rivoisy C, Tubach F, Roy C, Nicolas N, Mariette X, Salmon D, Lortholary O, Bourgarit A ; RATIO Group : Paradoxical anti-TNF-associated TB worsening : frequency and factors associated with IRIS. *Joint Bone Spine* 83 : 173-178, 2016
 83. Matsumoto T, Tanaka T : Continuation of anti-TNF therapy for rheumatoid arthritis in patients with active tuberculosis reactivated during anti-TNF medication is more beneficial than its cessation. *Infectious Diseases and Therapeutics* 3 : 35-37, 2015
 84. Winthrop KL, Chang E, Yamashita S, Iademarco MF, LoBue PA : Nontuberculous mycobacteria infections and anti-tumor necrosis factor-alpha therapy. *Emerg Infect Dis* 15 : 1556-1561, 2009
 85. Takazono T, Nakamura S, Imamura Y, Miyazaki T, Izumikawa K, Kakeya H, Yanagihara K, Kohno S : Paradoxical response to disseminated non-tuberculosis mycobacteriosis treatment in a patient receiving tumor necrosis factor- α inhibitor : a case report. *BMC Infect Dis* 14 : 114, 2014, doi : 10.1186/1471-2334-14-114.
 86. Lemoine M, Laurent C, Hanoy M, Leporrier J, François A, Guerrot D, Godin M : Immune Reconstitution Inflammatory Syndrome Secondary to *Mycobacterium kansasii* infection in a kidney transplant recipient. *Am J Transplant* 15 : 3255-3258, 2015
 87. Daum S, Moos V, Loddenkemper C, Schürmann C, Märker-Hermann E, Stein H, Zeitz M, Schneider T : Immune reconstitution inflammatory syndrome (IRIS) of the small bowel in an immunocompromised patient suffering from systemic lupus erythematosus and non-tuberculous mycobacteriosis. *Z Rheumatol* 67 : 277-278, 2008 (in German)
 88. Langan EA, Graetz V, Allerheiligen J, Zillikens D, Rupp J, Terheyden P : Immune checkpoint inhibitors and tuberculosis : an old disease in a new context. *Lancet Oncol* 21 : e55-e65, 2020
 89. Cadranet J, Canellas A, Matton L, Darrason M, Parrot A, Naccache JM, Lavolé A, Ruppert AM, Fallet V : Pulmonary complications of immune checkpoint inhibitors in patients with nonsmall cell lung cancer. *Eur Respir Rev* 28 : 190058, 2019, doi : 10.1183/16000617.0058-2019
 90. Takata S, Koh G, Han Y, Yoshida H, Shiroyama T, Takada H, Masuhiro K, Nasu S, Morita S, Tanaka A, Hashimoto S, Uriu K, Suzuki H, Tamura Y, Okamoto N, Nagai T, Hirashima T : Paradoxical response in a patient with non-small cell lung cancer who received nivolumab followed by anti-*Mycobacterium tuberculosis* agents. *J Infect Chemother* 25 : 54-58, 2019
 91. Fujita K, Yamamoto Y, Kanai O, Okamura M, Nakatani K, Mio T : Development of *Mycobacterium avium* Complex Lung Disease in Patients With Lung Cancer on Immune Checkpoint Inhibitors. *Open Forum Infect Dis* 7 : ofaa067, 2020 doi : 10.1093/ofid/ofaa067
 92. Ishii S, Tamiya A, Taniguchi Y, Tanaka T, Abe Y, Isa SI, Tsuyuguchi K, Suzuki K, Atagi S : Improvement of *Mycobacterium abscessus* pulmonary disease after nivolumab administration in a patient with advanced non-small cell lung cancer. *Intern Med* 57 : 3625-3629, 2018
 93. Cheng VC, Ho PL, Lee RA, Chan KS, Chan KK, Woo PC, Lau SK, Yuen KY : Clinical spectrum of paradoxical deterioration during antituberculosis therapy in non-HIV-infected patients. *Eur J Clin Microbiol Infect Dis* 21 : 803-809, 2002
 94. Breen RA, Smith CJ, Bettinson H, Dart S, Bannister B, Johnson MA, Lipman MC : Paradoxical reactions during tuberculosis treatment in patients with and without HIV co-infection. *Thorax* 59 : 704-707, 2004
 95. Yano S, Kobayashi K, Kato K, Tokuda Y, Ikeda T, Takeyama H : Paradoxical worsening of pulmonary *Mycobacterium abscessus*. *Respir Med* 101 : 868-870, 2007
 96. Shirayama R, Hamada K, Hayashi H, Tomoda K, Nakaya M, Yoshikawa M, Yoneda T, Narita N : Atypical mycobacteriosis (*Mycobacterium xenopi*) with "initial aggravation". *Nihon Kyobu Shikkan Gakkai Zasshi* 34 : 1035-9, 1996 (in Japanese)
 97. Singh N, Perfect JR : Immune reconstitution syndrome and exacerbation of infections after pregnancy. *Clin Infect Dis* 45 : 1192-1199, 2007
 98. Zenner D, Kruijshaar ME, Andrews N, Abubakar I : Risk of tuberculosis in pregnancy : a national, primary care-based cohort and self-controlled case series study. *Am J Respir Crit Care Med* 185 : 779-784, 2012

99. Cheng VC, Woo PC, Lau SK, Cheung CH, Yung RW, Yam LY, Yuen KY : Peripartum tuberculosis as a form of immunorestitution disease. *Eur J Clin Microbiol Infect Dis* 22 : 313-317, 2003
100. Shinohara T, Kagawa K, Okano Y, Sawada T, Kobayashi T, Takikawa M, Iwahara Y, Ogushi F : Disseminated tuberculosis after pregnancy progressed to paradoxical response to the treatment : report of two cases. *BMC Infect Dis* 16 : 284, 2016, doi : 10.1186/s12879-016-1624-x.
101. Tsao SM, Liu KS, Liao HH, Huang TL, Shen GH, Tsao TC, Lee YT : The clinical management of cesarean section-acquired *Mycobacterium abscessus* surgical site infections. *J Infect Dev Ctries* 8 : 184-192, 2014
102. Wu TS, Yang CH, Brown-Elliott BA, Chao AS, Leu HS, Wu TL, Lin CS, Griffith DE, Chiu CH : Postcesarean section wound infection caused by *Mycobacterium massiliense*. *J Microbiol Immunol Infect* 49 : 955-961, 2016