






ORIGINAL RESEARCH

Effect of Underlying Cardiometabolic Diseases on the Association Between Sedentary Time and All-Cause Mortality in a Large Japanese Population: A Cohort Analysis Based on the J-MICC Study

Teruhide Koyama , PhD; Etsuko Ozaki, PhD; Nagato Kuriyama, MD, PhD; Satomi Tomida, MD; Tamami Yoshida , MS; Ritei Uehara, MD, PhD; Keitaro Tanaka, MD, PhD; Megumi Hara, MD, PhD; Asahi Hishida, MD, PhD; Rieko Okada , MD, PhD; Yoko Kubo, BA; Isao Oze, MD, PhD; Yuriko N. Koyanagi, MD, PhD; Haruo Mikami, MD; Yohko Nakamura, PhD; Ippei Shimoshikiryo, PT, MS; Toshiro Takezaki, MD, DMSc; Sadao Suzuki, MD, PhD, MSc; Takahiro Otani, PhD; Kiyonori Kuriki, PhD, MSc; Naoyuki Takashima, MD, PhD; Aya Kadota , MD, PhD; Kokichi Arisawa, MD, PhD, MSc; Sakurako Katsuura-Kamano, PhD; Hiroaki Ikezaki , MD, PhD; Masayuki Murata, MD, PhD; Kenji Takeuchi, PhD, DDS; Kenji Wakai, MD, PhD; for the Japan Multi-Institutional Collaborative Cohort (J-MICC) Study Group*

BACKGROUND: This study aimed to determine the association between sedentary time and mortality with regard to leisure-time physical activity with or without cardiometabolic diseases such as hypertension, dyslipidemia, and diabetes mellitus.

METHODS AND RESULTS: Using data from the J-MICC (Japan Multi-Institutional Collaborative Cohort) Study, 64 456 participants (29 022 men, 35 434 women) were analyzed. Hazard ratios (HRs) and 95% CIs were used to characterize the relative risk of all-cause mortality to evaluate its association with sedentary time (categorical variables: <5, 5 to <7, 7 to <9, ≥9 h/d and 2-hour increments in exposure) according to the self-reported hypertension, dyslipidemia, and diabetes mellitus using a Cox proportional hazards model. A total of 2257 participants died during 7.7 years of follow-up. The corresponding HRs for each 2-hour increment in sedentary time among participants with all factors, no factors, hypertension, dyslipidemia, and diabetes mellitus were 1.153 (95% CI, 1.114–1.194), 1.125 (95% CI, 1.074–1.179), 1.202 (95% CI, 1.129–1.279), 1.176 (95% CI, 1.087–1.273), and 1.272 (95% CI, 1.159–1.396), respectively. Furthermore, when analyzed according to the combined different factors (hypertension, dyslipidemia, and diabetes mellitus), HRs increased with each additional factor, and participants reporting all 3 conditions had the highest HR of 1.417 (95% CI, 1.162–1.728) independently of leisure-time metabolic equivalents.

CONCLUSIONS: The association between sedentary time and increased mortality is stronger among patients with hypertension, dyslipidemia, and diabetes mellitus regardless of leisure-time physical activity in a large Japanese population.

Key Words: all-cause mortality ■ diabetes mellitus ■ dyslipidemia ■ hypertension ■ sedentary time

Correspondence to: Teruhide Koyama, PhD, Department of Epidemiology for Community Health and Medicine, Kyoto Prefectural University of Medicine, 465 Kajii-cho, Kamigyo-ku, Kyoto 602-8566, Japan. E-mail: tkoyama@koto.kpu-m.ac.jp

*A complete list of the Japan Multi-institutional Collaborative Cohort (J-MICC) Study Group can be found in the Appendix at the end of the article.

Supplementary Material for this article is available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.120.018293>

For Sources of Funding and Disclosures, see page 8.

© 2021 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

JAHA is available at: www.ahajournals.org/journal/jaha

CLINICAL PERSPECTIVE

What Is New?

- The association between sedentary time and increased mortality is stronger among participants with cardiometabolic diseases related to hypertension, dyslipidemia, and diabetes mellitus regardless of leisure-time metabolic equivalents in a large population-based cohort study of Japanese adults.

What Are the Clinical Implications?

- The association between sedentary time and the number of hypertension/dyslipidemia/diabetes mellitus traits increases the risk of mortality.
- The association between sedentary time and increased mortality persists regardless of leisure-time physical activity.
- Greater health literacy is needed to interrupt and reduce sedentary time in adults to promote preventive medicine.

Nonstandard Abbreviations and Acronyms

J-MICC	Japan Multi-Institutional Collaborative Cohort
LT-METS	leisure-time metabolic equivalents

Physical inactivity is recognized as a pandemic that requires global attention. Physical inactivity and its harmful effects on health is highly prevalent worldwide. The evidence of the effectiveness of physical activity promotion strategies make this problem a global public health priority.¹ Physical inactivity is also significant from the perspective of economic burden.² Some systematic reviews and meta-analyses on physical inactivity reported that sedentary behavior (sitting or reclining posture) is associated with negative health consequences³ including increased cardiovascular-specific, cancer-specific, and overall mortality.^{4–6} Although modest amounts of activity provide substantial benefits for postponing mortality,⁷ it is unclear whether leisure-time physical activity offsets the health risks of sedentary behavior.

Recently, we reported the association of sedentary time with cardiometabolic diseases such as hypertension, dyslipidemia, and diabetes mellitus in a large Japanese population.⁸ However, there are no studies on the association between sedentary time and mortality in patients with cardiometabolic diseases such as hypertension, dyslipidemia, and diabetes mellitus.

Previous studies have reported that sedentary time affects cardiovascular diseases⁹ and is associated with mortality.^{4,10,11} In sum, the hypothesis holds that longer sedentary times tend to increase the risk of death in patients with cardiometabolic diseases. On the other hand, although the critical mechanism is unknown, several studies have shown that sedentary behavior–related health disorders are independent of physical activity.^{12,13} Actually, our previous study showed sedentary time was associated with cardiometabolic diseases independent of leisure-time physical activity.⁸ This study therefore aimed to determine the association between sedentary time and mortality with regard to leisure-time physical activity with or without underlying diseases such as hypertension, dyslipidemia, and diabetes mellitus in a large Japanese population.

METHODS

Requests to access the data set should be consulted with the J-MICC (Japan Multi-Institutional Collaborative Cohort) Study central office via the corresponding author of this article.

Study Participants

In the present study, we evaluated participant data collected during the J-MICC Study.¹⁴ The cohort study collected and analyzed genetic and clinical data from the general Japanese population to detect and confirm gene–environment interactions related to lifestyle-associated diseases. The study participants were aged 35 to 69 years and were enrolled upon responding to study announcements, attending health checkup examinations commissioned by their local governments, visiting local health checkup centers, or visiting a cancer hospital from February 2004 through March 2014.

When there were no data on participants' daily activities including sitting time, the research sites were excluded. Then, we excluded participants who lacked the following data in the self-administered questionnaire: history of hypertension, dyslipidemia, or diabetes mellitus; smoking and drinking status; physical activity including sitting time; the medical history of ischemic heart disease and stroke; drug treatment for hypertension, dyslipidemia, or diabetes mellitus; and those followed up for <1 year, which was too short of a period to assess the daily impact of reduced physical activity caused by the disease.

Changes in residence status, including vital status, were confirmed from 2004 using the residential registry. Mortality data were obtained from the Ministry of Health, Labor, and Welfare of Japan, and the underlying causes of death were coded according to the

International Classification of Diseases, Tenth Revision (ICD-10). Registration of death was required by the Family Registration Law and was believed to be complete, except for subjects who died after moving from their original study area. The participants were followed up through December 2016 in 10 study areas or December 2017 in 4 areas, except for parts of 1 area (December 2012 or December 2013). Selection bias was avoided because sufficient follow-up was conducted by the research sites.

All study participants provided written informed consent. The study protocol was approved by the ethics committees at the Nagoya University Graduate School of Medicine (IRB No. 2010-939) and other institutions participating in the J-MICC Study. This study was conducted according to the principles enshrined in the World Medical Association Declaration of Helsinki.

Self-Administered Questionnaire Data

In this study, we evaluated data on clinical and lifestyle variables (smoking and drinking status, and physical activity including sitting time) obtained through self-administered questionnaires. Physical activity was determined using a format similar to that of the International Physical Activity Questionnaire.¹⁵ Leisure-time physical activity was measured in terms of leisure-time metabolic equivalents (LT-METs), as previously reported.^{16,17} In brief, metabolic equivalent hours per day of leisure-time activity were estimated by multiplying the reported daily time spent in each activity by its corresponding metabolic equivalent intensity. We divided the LT-METs groups into quartiles (Q1–Q4). The duration of sitting time was asked with 8 possible responses: none, <1, 1 to <3, 3 to <5, 5 to <7, 7 to <9, 9 to <11, and ≥11 h/d. Sitting time was then categorized on the basis of the quartile value as mentioned in a previous study⁸: <5, 5 to <7, 7 to <9, or ≥9 h/d. Medical history and medication history were assessed using self-administered questionnaires. Hypertension, dyslipidemia (specifically hyperlipidemia), and diabetes mellitus were defined by the presence or absence of medical history and/or current use of medication.

Statistical Analysis

Continuous variables were expressed as means and SDs, and categorical variables were expressed as frequencies and percentages. Intergroup comparisons were performed using 1-way ANOVA for continuous variables and χ^2 tests for categorical variables. Person-years of follow-up were calculated from the date of filling out the baseline questionnaire to the time of death, moving out of the area, or the end of the follow-up, whichever came first. Hazard ratios (HRs) and 95% CIs

were used to characterize the relative risk of all-cause mortality to evaluate its association with sedentary time (<5 hours was used as a reference; categorical variables: <5, 5 to <7, 7 to <9, ≥9 and 2-hour increments in exposure) according to the self-reported hypertension, dyslipidemia, and diabetes mellitus using a Cox proportional hazards model. The model was adjusted for age (operationalized as a continuous variable), sex, research area, LT-METs (Q1–Q4), drinking and smoking status (never, former, and current), and history of ischemic heart disease and stroke, and history of medication for hypertension, dyslipidemia, and diabetes mellitus. The incidence of mortality was calculated using the Kaplan-Meier method. Tests for linear trends (eg, *P* trend tests) were conducted by including 4 groups according to LT-METs (Q1–Q4) as ordinal variables. Using G*Power (<http://www.gpower.hhu.de>), we verified that the sample size was sufficient. A post hoc power calculation showed the sample size in this study achieved a power of 0.80 at 0.05 α level for a 2-sided test. All statistical analyses were performed using SPSS software version 25 (IBM Japan, Tokyo, Japan) and JMP 13 software (SAS Institute, Cary, NC), and *P*<0.005 was considered to indicate statistical significance.

RESULTS

Figure 1 shows the study participant flowchart. Of the 14 research sites, 2 did not collect data on participants' daily activities including sitting time. Excluding the participants from these 2 research sites, 72 712 participants were initially included in the current study (data set version 20200312). We verified the reproducibility of the analysis results and confirmed that the results were correct.

Participants' characteristics, including drinking and smoking status, medical history of hypertension/dyslipidemia/diabetes mellitus, and distribution of age and sex according to sedentary time are presented in Table 1. Of 64 456 participants included in the current analysis, 24 304 (37.7%), 14 596 (22.6%), 10 481 (16.3%), and 15 075 (23.4%) spent <5, 5 to <7, 7 to <9, and ≥9 hours per day being sedentary, respectively. A total of 2257 participants died during the 7.7 years (mean) of follow-up. Mortality increased with longer sedentary time, and the mortality rate (cases per 1000 person-years) increased from 3.93 in the <5-hours group to 6.26 in the ≥9-hours group.

Results from the multivariate model (ie, adjusted HR [95% CI]) of all-cause mortality according to medical history of hypertension, dyslipidemia, and diabetes mellitus are shown in Table 2. Compared with participants who spent <5 hours of sedentary time, those who had longer sedentary time tended to have significantly higher HRs, especially in the ≥9-hours group.




Figure 1. Flowchart of the study participants.
J-MICC indicates Japan Multi-Institutional Collaborative Cohort.

The corresponding HR for each 2-hour increment in sedentary time was 1.153 (95% CI, 1.114–1.194) among all participants. The corresponding HRs for

each 2-hour increment in sedentary time among participants with no factors, hypertension, dyslipidemia, and diabetes mellitus were 1.125 (95% CI, 1.074–1.179),

Table 1. Characteristics of Participants According to Sedentary Time

Sedentary Time	<5 h	5 to <7 h	7 to <9 h	≥9 h
	n=24 304	n=14 596	n=10 481	n=15 075
Age, y	54.7±9.45	55.4±9.43	54.8±9.55	53.4±9.64
Sex, men	9910 (40.8)	5719 (39.2)	4678 (44.6)	8715 (57.8)
No. of deaths	789 (3.2)	449 (3.1)	366 (3.5)	653 (4.3)
Person-years	200 658	114 237	79 389	104 386
Mortality rate (per 1000 person-years)	3.93	3.93	4.61	6.26
Drinking status				
Current	13 261 (54.6)	8003 (54.8)	6022 (57.5)	9318 (61.8)
Former	492 (2.0)	352 (2.4)	249 (2.4)	465 (3.1)
Never	10 551 (43.4)	6241 (42.8)	4210 (40.2)	5292 (35.1)
Smoking status				
Current	4226 (17.4)	2157 (14.8)	1606 (15.3)	2885 (19.1)
Former	4990 (20.5)	3115 (21.3)	2537 (24.2)	4394 (29.1)
Never	15 088 (62.1)	9324 (63.9)	6338 (60.5)	7796 (51.7)
Hypertension	4697 (19.3)	3007 (20.6)	2178 (20.8)	3078 (20.4)
Dyslipidemia	4010 (16.5)	2928 (20.1)	2123 (20.3)	3124 (20.7)
Diabetes mellitus	1380 (5.7)	845 (5.8)	659 (6.3)	1005 (6.7)
Stroke	366 (1.5)	227 (1.6)	159 (1.5)	258 (1.7)
IHD	590 (2.4)	366 (2.5)	316 (3.0)	471 (3.1)
LT-METs, h/d	2.21±3.54	2.32±3.23	2.06±2.89	1.71±2.52

Data are presented as mean±SD or number (percentage). IHD indicates ischemic heart disease; and LT-METs, leisure-time metabolic equivalents.

1.202 (95% CI, 1.129–1.279), 1.176 (95% CI, 1.087–1.273), and 1.272 (95% CI, 1.159–1.396), respectively. Furthermore, when analyzed according to the combined different factors (hypertension, dyslipidemia, and diabetes mellitus), HRs increased with each additional factor, and participants reporting all 3 conditions had the highest HR of 1.417 (95% CI, 1.162–1.728). The Kaplan-Meier cumulative mortality data stratified by group are shown in Figure 2. The risk of mortality significantly increased among the group with a longer sedentary time ($P<0.001$ using the log-rank test).

We also reported results of HRs for each 2-hour increment in sedentary time according to the LT-METs quartile. Table S1 shows the characteristics of participants in each LT-METs quartile. As the LT-METs quartile increased, the mean age and proportion of participants reporting hypertension/dyslipidemia/diabetes mellitus increased; however, mortality rate (cases per 1000 person-years) decreased with higher LT-METs from 5.14 in the Q1 group to 4.19 in the Q4 group. In the all- and no-factors groups, participants with higher LT-METs tended to have significantly lower HRs; particularly in the fourth quartile of LT-METs (Q4), the HR for the no-factors group was not significant (HR, 1.091; 95% CI, 0.982–1.211) (Table 3). However, most HRs of sedentary time and mortality remained statistically significant, regardless of the LT-METs group.

DISCUSSION

Considerable evidence suggests that sedentary behavior is associated with increased mortality.^{13,18–20} The results of this study confirm the association that each 2-hour increment in sedentary time was associated with a 15.3% increase in the risk of mortality among all participants and a 41.7% increase in the risk of mortality among participants with hypertension, dyslipidemia, and diabetes mellitus. Thus, our results suggest that the well-established association between sedentary time and increased mortality is stronger among patients with a medical history of hypertension, dyslipidemia, and diabetes mellitus. In general, common correlates of sedentary behavior are known to differ by ethnicity.²¹ Few studies have examined the association between sedentary time and mortality in Japan. The only 2 studies that showed an association between sedentary time and mortality involved limited situations: television viewing time and mortality from stroke and coronary artery disease,²² and occupational sitting time and all-cause mortality.²³ To the best of our knowledge, this is the first study to determine that each 2-hour increment in sedentary time and the number of hypertension/dyslipidemia/diabetes mellitus traits increases the risk of mortality in Japanese adults.

Table 2. Associations Between Death and Sedentary Time According to Self-Reported Cardiometabolic Disease

	No.	No. of Deaths	Sedentary Time												2-h Increments in Sedentary Time	
			<5 h			5 to <7 h			7 to <9 h			≥9 h			HR	95% CI
			Reference	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI			
All	64 456	2257	1.000	1.031	0.918–1.158	1.205	1.064–1.364	1.540	1.386–1.712	1.153	1.114–1.194	1.000	1.000	1.000	1.000	
Hypertension	12 960	702	1.000	1.075	0.868–1.329	1.328	1.063–1.659	1.729	1.430–2.090	1.202	1.129–1.279	1.000	1.000	1.000	1.000	
Dyslipidemia	12 185	454	1.000	1.133	0.871–1.453	1.405	1.069–1.847	1.608	1.259–2.055	1.176	1.087–1.273	1.000	1.000	1.000	1.000	
Diabetes mellitus	3889	306	1.000	1.566	1.130–2.170	1.728	1.225–2.438	2.142	1.594–2.877	1.272	1.159–1.396	1.000	1.000	1.000	1.000	
No. of hypertension/dyslipidemia/diabetes mellitus																
None	42 911	1231	1.000	1.017	0.871–1.188	1.129	0.953–1.339	1.447	1.254–1.668	1.125	1.074–1.179	1.000	1.000	1.000	1.000	
1	15 067	666	1.000	0.896	0.719–1.117	1.214	0.968–1.522	1.619	1.336–1.962	1.181	1.108–1.259	1.000	1.000	1.000	1.000	
2	5467	284	1.000	1.311	0.943–1.823	1.386	0.967–1.987	1.741	1.282–2.366	1.192	1.081–1.316	1.000	1.000	1.000	1.000	
3	1011	76	1.000	2.269	1.112–4.630	2.684	1.324–5.437	3.175	1.624–6.210	1.417	1.162–1.728	1.000	1.000	1.000	1.000	

Adjusted for age, sex, research area, leisure-time-metabolic equivalents, drinking and smoking status, ischemic heart disease, stroke, and history of medication for hypertension, dyslipidemia, and diabetes mellitus. HR indicates hazard ratio.

In this study, the population characteristics varied across LT-METs quartiles, as shown in Table S1. As the LT-METs quartile increased, the mean age and the proportion of participants with hypertension and/or dyslipidemia and/or diabetes mellitus also increased. Interestingly, the number of deaths decreased with increasing LT-METs quartile. It was speculated that in the higher-quartile LT-METs group, factors such as being elderly and having underlying diseases might have affected health consciousness. In general, the strength of the associations between motivational regulations and physical activity behavior varied across age groups.^{24,25} Health promotion involving physical activity is well established, and researchers have sought to explore the reasons why some people are physically active, whereas others are not.^{26–29} Older individuals exhibit greater concerns about health outcomes, thus motivating them to perform physical activity.³⁰ It is reasonable to expect that older adults would show more concern for physical and psychological health issues when making their physical activity decisions.³¹ Based on these results, the higher quartile LT-METs group included older subjects and participants with underlying diseases, which may have affected the association of mortality with sedentary time. On the other hand, it is suggested that increasing physical activity during leisure time after the onset of a disease or because of increasing age may be considered a part of secondary or tertiary prevention of the disease, but it is not sufficient for primary prevention. This highlights the necessity of an intervention for the allocation of daily physical activity behaviors among people who do not have a disease and who lack health awareness, for the purpose of promoting preventive medicine. Sedentary time was associated with cardiometabolic diseases such as hypertension, dyslipidemia, and diabetes mellitus among Japanese population.⁸ In sum, reducing sedentary time may be an effective measure to prevent cardiometabolic diseases and death. However, some people cannot change their behavior during working hours. Because sedentary behavior accounts for 55% to 60% of the total duration of daytime activity,³² the use of leisure time on weekdays can reduce the duration of sedentary time during the day outside of working hours.

We examined the effect of factors (defined as self-reported hypertension, dyslipidemia, or diabetes mellitus in this study) on the relationship between sedentary time and risk of death. Sedentary behavior in combination with chronic diseases,³³ including hypertension³⁴ and diabetes mellitus,²⁰ increased all-cause mortality risk. Although the relationship between sedentary time and death among participants with dyslipidemia is often unknown, our results showed that participants with at least 1 factor had a higher risk of mortality than




Figure 2. Kaplan-Meier cumulative mortality classified by sedentary time (Group 1: <5 hours; Group 2: 5 to <7 hours; Group 3: 7 to <9 hours; and Group 4: ≥9 hours). *P*<0.001 using the log-rank test.

did participants with none. As inferred from previous reports, the association between sedentary time and diabetes mellitus has a stronger risk for mortality than another medical history.³⁵ Furthermore, increasing the number of people with hypertension/dyslipidemia/diabetes mellitus increases the risk of mortality (HR), which is as expected. Importantly, a strong association between sedentary time and mortality was observed among participants with hypertension/dyslipidemia/diabetes mellitus independently of LT-METs grouping; therefore, there is a need to increase awareness among healthy people who lack health literacy to interrupt and reduce their sedentary time.

Despite our novel findings, this study has some limitations. This study used a self-administered

questionnaire at baseline survey to evaluate sedentary time and medical history. Changes in risk factors since the baseline survey could not be taken into account, such as sitting time, exercise during leisure time, drinking, and smoking. Although a questionnaire that evaluates sitting time is controversial, the International Physical Activity Questionnaire is a widely accepted international physical activity surveillance instrument.³⁶ Self-reported measures continue to be the most widely used method for assessing these behaviors. However, evidence from one review suggests that single-item self-reported measures generally underestimate sedentary time compared with device-based measures.³⁷ The strength of this study is the inclusion of a large number of participants, which prevented sample bias

Table 3. Associations Between Death and 2-Hour Increments in Sedentary Time According to LT-METs Quartile

LT-METs	Q1		Q2		Q3		Q4		P Value for Trend
	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	
All	1.169	1.100–1.242	1.143	1.067–1.226	1.165	1.082–1.254	1.126	1.042–1.217	<0.001
Hypertension	1.257	1.123–1.406	1.187	1.048–1.344	1.162	1.021–1.322	1.177	1.022–1.356	<0.001
Dyslipidemia	1.090	0.945–1.257	1.230	1.045–1.448	1.212	1.022–1.438	1.206	1.026–1.417	<0.001
Diabetes mellitus	1.352	1.151–1.587	1.205	0.992–1.464	1.243	1.033–1.495	1.229	0.965–1.567	<0.001
No. of hypertension/dyslipidemia/diabetes mellitus									
None	1.147	1.058–1.243	1.111	1.012–1.220	1.147	1.035–1.270	1.091	0.982–1.211	<0.001
1	1.197	1.067–1.344	1.165	1.024–1.325	1.212	1.062–1.383	1.157	1.001–1.337	<0.001
2	1.192	0.995–1.428	1.364	1.114–1.671	1.072	0.869–1.323	1.149	0.931–1.418	<0.001
3	1.506	1.066–2.126	1.029	0.664–1.594	1.612	1.071–2.427	2.044	1.138–3.670	0.018

Adjusted for age, sex, research area, drinking and smoking status, ischemic heart disease, stroke and history of medication for hypertension, dyslipidemia, and diabetes mellitus. HR indicates hazard ratio; and LT-METs, leisure-time metabolic equivalents.

and implementation of a population-based cohort design.

CONCLUSIONS

This study showed that the association between sedentary time and increased mortality is stronger among patients with cardiometabolic diseases related to hypertension, dyslipidemia, and diabetes mellitus regardless of LT-METs in a large Japanese population.

APPENDIX

J-MICC Study Group: Kenji Wakai, Kenji Takeuchi, Haruo Mikami, Hiroki Nagase, Hiroto Narimatsu, Kiyonori Kuriki, Keitaro Matsuo, Sadao Suzuki, Asahi Hishida, Yoshikuni Kita, Katsuyuki Miura, Ritei Uehara, Kokichi Arisawa, Hiroaki Ikezaki, Keitaro Tanaka, and Toshiro Takezaki.

ARTICLE INFORMATION

Received June 30, 2020; accepted April 7, 2021.

Affiliations

Department of Epidemiology for Community Health and Medicine (T.K., E.O., N.K., S.T., T.Y., R.U.) and Department of Endocrine and Breast Surgery (S.T.), Kyoto Prefectural University of Medicine, Kyoto, Japan; Shizuoka Graduate University of Public Health, Shizuoka, Japan (N.K.); Department of Preventive Medicine, Faculty of Medicine, Saga University, Saga, Japan (K. Tanaka, M.H.); Department of Preventive Medicine, Nagoya University Graduate School of Medicine, Aichi, Japan (A.H., R.O., Y.K., K. Takeuchi, K.W.); Division of Cancer Epidemiology and Prevention (I.O.) and Division of Cancer Information and Control (Y.N.K.), Aichi Cancer Center Research Institute, Aichi, Japan; Cancer Prevention Center, Chiba Cancer Center Research Institute, Chiba, Japan (H.M., Y.N.); Department of International Island and Community Medicine, Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima, Japan (I.S., T.T.); Department of Public Health, Nagoya City University Graduate School of Medical Sciences, Aichi, Japan (S.S., T.O.); Laboratory of Public Health, School of Food and Nutritional Sciences, University of Shizuoka, Japan (K.K.); Department of Public Health, Faculty of Medicine, Kindai University, Osaka, Japan (N.T.); Department of Public Health, Shiga University of Medical Science, Shiga, Japan (N.T., A.K.); Department of Preventive Medicine, Tokushima University Graduate School of Biomedical Sciences, Tokushima, Japan (K.A., S.K.-K.); Department of General Internal Medicine, Kyushu University Hospital, Fukuoka, Japan (H.I., M.M.); and Department of Comprehensive General Internal Medicine, Faculty of Medical Sciences, Kyushu University, Fukuoka, Japan (H.I.).

Acknowledgments

The authors thank Drs Hamajima and Tanaka for their work in initiating and organizing the J-MICC Study as former principal investigators.

Sources of Funding

This study was supported by Grants-in-Aid for Scientific Research for Priority Areas of Cancer (No. 17015018) and Innovative Areas (No. 221S0001) and by the Japan Society for the Promotion of Science (JSPS) KAKENHI Grant (No. 16H06277 [CoBiA]) from the Japanese Ministry of Education, Culture, Sports, Science and Technology.

Disclosures

None.

Supplementary Material

Table S1

REFERENCES

- Kohl HW III, Craig CL, Lambert EV, Inoue S, Alkandari JR, Leetongin G, Kahlmeier S. The pandemic of physical inactivity: global action for public health. *Lancet*. 2012;380:294–305. DOI: 10.1016/S0140-6736(12)60898-8.
- Ding D, Lawson KD, Kolbe-Alexander TL, Finkelstein EA, Katzmarzyk PT, van Mechelen W, Pratt M. The economic burden of physical inactivity: a global analysis of major non-communicable diseases. *Lancet*. 2016;388:1311–1324. DOI: 10.1016/S0140-6736(16)30383-X.
- de Rezende LF, Rey-López JP, Matsudo VK, do Carmo Luiz O. Sedentary behavior and health outcomes among older adults: a systematic review. *BMC Public Health*. 2014;14:333. DOI: 10.1186/1471-2458-14-333.
- Ekelund U, Brown WJ, Steene-Johannessen J, Fagerland MW, Owen N, Powell KE, Bauman AE, Lee IM. Do the associations of sedentary behaviour with cardiovascular disease mortality and cancer mortality differ by physical activity level? A systematic review and harmonised meta-analysis of data from 850 060 participants. *Br J Sports Med*. 2019;53:886–894.
- Ekelund U, Tarp J, Steene-Johannessen J, Hansen BH, Jefferis B, Fagerland MW, Whincup P, Diaz KM, Hooker SP, Chernofsky A, et al. Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: systematic review and harmonised meta-analysis. *BMJ*. 2019;366:i4570. DOI: 10.1136/bmj.i4570.
- Rojer AGM, Ramsey KA, Trappenburg MC, van Rijnssen NM, Otten RHJ, Heymans MW, Pijnappels M, Meskers CGM, Maier AB. Instrumented measures of sedentary behaviour and physical activity are associated with mortality in community-dwelling older adults: a systematic review, meta-analysis and meta-regression analysis. *Ageing Res Rev*. 2020;61:101061. DOI: 10.1016/j.arr.2020.101061.
- Arem H, Moore SC, Patel A, Hartge P, Berrington de Gonzalez A, Viswanathan K, Campbell PT, Freedman M, Weiderpass E, Adami HO, et al. Leisure time physical activity and mortality: a detailed pooled analysis of the dose-response relationship. *JAMA Intern Med*. 2015;175:959–967. DOI: 10.1001/jamainternmed.2015.0533.
- Koyama T, Kuriyama N, Ozaki E, Tomida S, Uehara R, Nishida Y, Shimano C, Hishida A, Tamura T, Tsukamoto M, et al. Sedentary time is associated with cardiometabolic diseases in a large Japanese population: a cross-sectional study. *J Atheroscler Thromb*. 2020;27:1097–1107. DOI: 10.5551/jat.54320.
- Young DR, Hivert M-F, Alhassan S, Camhi SM, Ferguson JF, Katzmarzyk PT, Lewis CE, Owen N, Perry CK, Siddique J, et al. Sedentary behavior and cardiovascular morbidity and mortality: a science advisory from the American Heart Association. *Circulation*. 2016;134:e262–e279. DOI: 10.1161/CIR.0000000000000440.
- Garcia JM, Duran AT, Schwartz JE, Booth JN III, Hooker SP, Willey JZ, Cheung YK, Park C, Williams SK, Sims M, et al. Types of sedentary behavior and risk of cardiovascular events and mortality in blacks: the Jackson Heart Study. *J Am Heart Assoc*. 2019;8:e010406. DOI: 10.1161/JAHA.118.010406.
- Evenson KR, Wen F, Herring AH. Associations of accelerometry-assessed and self-reported physical activity and sedentary behavior with all-cause and cardiovascular mortality among US adults. *Am J Epidemiol*. 2016;184:621–632.
- Loprinzi PD, Frith E. Effects of sedentary behavior, physical activity, frequency of protein consumption, lower extremity strength and lean mass on all-cause mortality. *J Lifestyle Med*. 2018;8:8–15. DOI: 10.15280/jlm.2018.8.1.8.
- Ekelund U, Steene-Johannessen J, Brown WJ, Fagerland MW, Owen N, Powell KE, Bauman A, Lee IM. Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women. *Lancet*. 2016;388:1302–1310. DOI: 10.1016/S0140-6736(16)30370-1.
- Hamajima N. The Japan Multi-Institutional Collaborative Cohort Study (J-MICC Study) to detect gene-environment interactions for cancer. *Asian Pac J Cancer Prev*. 2007;8:317–323.
- Craig CL, Marshall AL, Sjoström M, Bauman AE, Booth ML, Ainsworth BE, Pratt M, Ekelund U, Yngve A, Sallis JF, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc*. 2003;35:1381–1395. DOI: 10.1249/01.MSS.0000078924.61453.FB.
- Hara M, Higaki Y, Taguchi N, Shinchi K, Morita E, Naito M, Hamajima N, Takashima N, Suzuki S, Nakamura A, et al. Effect of the PPARG2

- Pro12Ala polymorphism and clinical risk factors for diabetes mellitus on HbA1c in the Japanese general population. *J Epidemiol.* 2012;22:523–531. DOI: 10.2188/jea.JE20120078.
17. Hara M, Hachiya T, Sutoh Y, Matsuo K, Nishida Y, Shimano C, Tanaka K, Shimizu A, Ohnaka K, Kawaguchi T, et al. Genomewide association study of leisure-time exercise behavior in Japanese adults. *Med Sci Sports Exerc.* 2018;50:2433–2441. DOI: 10.1249/MSS.00000000000001712.
 18. Patterson R, McNamara E, Tainio M, de Sa TH, Smith AD, Sharp SJ, Edwards P, Woodcock J, Brage S, Wijndaele K. Sedentary behaviour and risk of all-cause, cardiovascular and cancer mortality, and incident type 2 diabetes: a systematic review and dose response meta-analysis. *Eur J Epidemiol.* 2018;33:811–829.
 19. Powell C, Herring MP, Dowd KP, Donnelly AE, Carson BP. The cross-sectional associations between objectively measured sedentary time and cardiometabolic health markers in adults—a systematic review with meta-analysis component. *Obes Rev.* 2018;19:381–395.
 20. Wilmot EG, Edwardson CL, Achana FA, Davies MJ, Gorely T, Gray LJ, Khunti K, Yates T, Biddle SJ. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. *Diabetologia.* 2012;55:2895–2905. DOI: 10.1007/s00125-012-2677-z.
 21. Biddle GJH, Edwardson CL, Rowlands AV, Davies MJ, Bodicoat DH, Hardeman W, Eborall H, Sutton S, Griffin S, Khunti K, et al. Differences in objectively measured physical activity and sedentary behaviour between white Europeans and South Asians recruited from primary care: cross-sectional analysis of the PROPELS trial. *BMC Public Health.* 2019;19:95. DOI: 10.1186/s12889-018-6341-5.
 22. Ikehara S, Iso H, Wada Y, Tanabe N, Watanabe Y, Kikuchi S, Tamakoshi A. Television viewing time and mortality from stroke and coronary artery disease among Japanese men and women—the Japan Collaborative Cohort Study. *Circ J.* 2015;79:2389–2395.
 23. Kikuchi H, Inoue S, Odagiri Y, Inoue M, Sawada N, Tsugane S. Occupational sitting time and risk of all-cause mortality among Japanese workers. *Scand J Work Environ Health.* 2015;41:519–528.
 24. Brunet J, Sabiston CM. Exploring motivation for physical activity across the adult lifespan. *Psychol Sport Exerc.* 2011;12:99–105. DOI: 10.1016/j.psychsport.2010.09.006.
 25. Egli T, Bland HW, Melton BF, Czech DR. Influence of age, sex, and race on college students' exercise motivation of physical activity. *J Am Coll Health.* 2011;59:399–406. DOI: 10.1080/07448481.2010.513074.
 26. Franco MR, Tong A, Howard K, Sherrington C, Ferreira PH, Pinto RZ, Ferreira ML. Older people's perspectives on participation in physical activity: a systematic review and thematic synthesis of qualitative literature. *Br J Sports Med.* 2015;49:1268–1276. DOI: 10.1136/bjsports-2014-094015.
 27. Lee AM, Chavez S, Bian J, Thompson LA, Gurka MJ, Williamson VG, Modave F. Efficacy and effectiveness of mobile health technologies for facilitating physical activity in adolescents: scoping review. *JMIR Mhealth Uhealth.* 2019;7:e11847. DOI: 10.2196/11847.
 28. O'Halloran PD, Blackstock F, Shields N, Holland A, Iles R, Kingsley M, Bernhardt J, Lannin N, Morris ME, Taylor NF. Motivational interviewing to increase physical activity in people with chronic health conditions: a systematic review and meta-analysis. *Clin Rehabil.* 2014;28:1159–1171. DOI: 10.1177/0269215514536210.
 29. Zubala A, MacGillivray S, Frost H, Kroll T, Skelton DA, Gavine A, Gray NM, Toma M, Morris J. Promotion of physical activity interventions for community dwelling older adults: a systematic review of reviews. *PLoS One.* 2017;12:e0180902. DOI: 10.1371/journal.pone.0180902.
 30. Trujillo KM, Brougham RR, Walsh DA. Age differences in reasons for exercising. *Curr Psychol.* 2004;22:348–367. DOI: 10.1007/s12144-004-1040-z.
 31. Molanorouzi K, Khoo S, Morris T. Motives for adult participation in physical activity: type of activity, age, and gender. *BMC Public Health.* 2015;15:66. DOI: 10.1186/s12889-015-1429-7.
 32. Dunstan DW, Howard B, Healy GN, Owen N. Too much sitting—a health hazard. *Diabetes Res Clin Pract.* 2012;97:368–376.
 33. Zhao R, Bu W, Chen Y, Chen X. The dose-response associations of sedentary time with chronic diseases and the risk for all-cause mortality affected by different health status: a systematic review and meta-analysis. *J Nutr Health Aging.* 2020;24:63–70. DOI: 10.1007/s12603-019-1298-3.
 34. Joseph G, Marott JL, Torp-Pedersen C, Biering-Sørensen T, Nielsen G, Christensen AE, Johansen MB, Schnohr P, Sogaard P, Mogelvang R. Dose-response association between level of physical activity and mortality in normal, elevated, and high blood pressure. *Hypertension.* 2019;74:1307–1315. DOI: 10.1161/HYPERTENSIONAHA.119.13786.
 35. Chastin SFM, De Craemer M, De Cocker K, Powell L, Van Cauwenberg J, Dall P, Hamer M, Stamatakis E. How does light-intensity physical activity associate with adult cardiometabolic health and mortality? Systematic review with meta-analysis of experimental and observational studies. *Br J Sports Med.* 2019;53:370–376.
 36. Bauman A, Bull F, Chey T, Craig CL, Ainsworth BE, Sallis JF, Bowles HR, Hagstromer M, Sjostrom M, Pratt M. The international prevalence study on physical activity: results from 20 countries. *Int J Behav Nutr Phys Act.* 2009;6:21.
 37. Prince SA, Cardilli L, Reed JL, Saunders TJ, Kite C, Douillette K, Fournier K, Buckley JP. A comparison of self-reported and device measured sedentary behaviour in adults: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act.* 2020;17:31.

SUPPLEMENTAL MATERIAL

Table S1. Characteristics of participants according to LT-METs quartile.

LT-METs	Q1 n = 17021		Q2 n = 16530		Q3 n = 14889		Q4 n = 16016		p-value	
Age (year)	52.5	±9.40	53.4	±9.35	55.5	±9.35	57.1	±9.32	<0.001	
Sex (men)	7065	41.5%	7177	43.4%	6925	46.5%	7855	49.0%	<0.001	
No. of death	669	3.9%	561	3.4%	514	3.5%	513	3.2%	0.003	
Person-year	130130		128965		117010		122565			
Mortality rate (/1000 person-years)	5.14		4.35		4.39		4.19			
Drinking status										
	Current	9011	52.9%	9292	56.2%	8617	57.9%	9684	60.5%	
	Former	475	2.8%	387	2.3%	349	2.3%	347	2.2%	<0.001
	Never	7535	44.3%	6851	41.4%	5923	39.8%	5985	37.4%	
Smoking status										
	Current	3759	22.1%	2673	16.2%	2298	15.4%	2144	13.4%	
	Former	3237	19.0%	3738	22.6%	3740	25.1%	4321	27.0%	<0.001
	Never	10025	58.9%	10119	61.2%	8851	59.4%	9551	59.6%	
Hypertension	3003		3152		3236		3569		<0.001	
Dyslipidemia	2616		3049		3099		3421		<0.001	
Diabetes	855		894		1001		1139		<0.001	
Stroke	230		242		242		296		0.002	
IHD	428		408		428		479		0.006	

Data are presented as mean (±SD), or number (percentage).

IHD, ischemic heart disease; LT-METs, Leisure-time-metabolic equivalents.