

Fertilization failure and oocyte activation

Shuji Yamano^{*}, Koji Nakagawa[†], Hisayo Nakasaka[†], and Toshihiro Aono[†]

^{*}Center for Maternity and Perinatal Care, and [†]Department of Obstetrics and Gynecology, The University of Tokushima School of Medicine, Tokushima, Japan

Abstract : Recent reports have suggested that when unfertilized oocytes with a spermatozoon after intracytoplasmic sperm injection (ICSI) are properly activated, the activated oocytes develop normally similar to fertilized oocytes. However, human oocytes do not easily respond to universal activators of mammalian oocytes like ethanol or calcium ionophore A23187, which induce a calcium transient in ooplasm. Puromycin activates human oocytes at a rate of 90%, but more than two thirds of the parthenogenones possess 2 haploid pronuclei or 1 diploid pronucleus without extrusion of the second polar body. Therefore, the activation method which produces one pronucleus with extrusion of the second polar body in oocytes without a spermatozoon is necessary for producing embryos with normal karyotypes. Recently, we found the oocyte activation method which produced parthenogenones displaying one pronucleus with extrusion of the second polar body. Using our method (a combination of calcium ionophore A23187 and puromycin), the activation rate was approximately 90% and the proportion of parthenogenones displaying one pronucleus with extrusion of the second polar body was approximately 80% in human aged and mouse young oocytes. When human unfertilized oocytes following ICSI were activated by this method, two pronuclei were formed with extrusion of the second polar body in 30% of the oocytes. Four cleaved parthenogenones (or embryos) showed normal karyotypes. However, the cytotoxic, teratogenic and mutagenetic activity of Ca ionophore and puromycin should be approved prior to the clinical adaptation of the method. *J. Med. Invest.* 47 : 1-8, 2000

Key words : fertilization, oocyte activation, parthenogenone, MPF, MAPK

INTRODUCTION

In-vitro fertilization (IVF) is a technique which offers a possibility for fertilization to infertile couples and has been used world-wide for aiding infertile couples with tubal blockage. However, the technique is not effective for severe oligozoospermic patients because few spermatozoa of the patients can penetrate the zona pellucida. For more than a decade, the fertilization failure of spermatozoa from oligozoospermic patients was one of the most important problems to be solved for improving the

pregnancy rate of oligozoospermic patients. The appearance of intra-cytoplasmic sperm injection (ICSI) dramatically improved the fertilization and pregnancy rates of oligozoospermic patients and ICSI showed that penetration of spermatozoa through the zona pellucida and fusion between a sperm cell membrane and oolemma are not essential for early development of embryos (1). However, fertilization failure still occurs in ICSI (1-3). Approximately 80% of unfertilized oocytes following ICSI contain a spermatozoon (4). This suggests that the process from meiosis to mitosis was impaired in the unfertilized oocytes and the fertilization failure is defined as failure of the M-G1 transition in metaphase II (MII) oocytes. The failure of the M-G1 transition may be repaired by artificial oocyte activation. In this review, we discuss the possible treatment for failure of the M-G1 transition in MII oocyte.

Received for publication December 24, 1999 ; accepted January 13, 2000.

Address correspondence and reprint requests to Shuji Yamano, M.D., Center for Maternity and Perinatal Care, The University of Tokushima School of Medicine, Kuramoto-cho, Tokushima 770-8503, Japan and Fax : +81-88-631-2630.

FAILURE OF OOCYTE ACTIVATION IN ICSI

In mammals, except dogs and foxes, oocytes which are ovulated from ovaries are arrested at M II and they contain a high level of maturation (or M-phase) promoting factor (MPF : the universal regulator of the G 2-M transition in eukaryotic cells, which consists of cdc2 kinase and cyclin B). Cytostatic factor (CSF), which stabilizes MPF activity is also kept high in MII oocytes (5). When one spermatozoon contacts the oolemma or penetrates into the ooplasm, calcium oscillation occurs (6-8). The mechanism that induces calcium oscillation in ooplasm remains unknown. However the success of ICSI strongly suggests that a factor which induces calcium oscillation in ooplasm (so called oscillin) is present in spermatozoa and is directly introduced into the ooplasm, not via a sperm receptor (4). Calmodulin-dependent kinase II (CaM KII) is activated by a Ca^{2+} transient and the activation of the kinase probably turns on the cyclin degradation machinery, resulting in cyclin B degradation and sister chromatid separation (9) (Fig.1). The process of M-G1 transition in MII oocytes is called " oocyte activation ". In in vitro fertilization (IVF) 5-32% of unfertilized oocytes have a spermatozoon in the ooplasm (10-15). This suggests that at least two-thirds of spermatozoa, which penetrate oocytes possesses an ability to activate MII oocytes.

ICSI is the most efficient and widely accepted treatment for aiding severe oligozoospermic patients, a group of azoospermic patients and patients who experienced previous fertilization failure using conventional IVF and embryo transfer (ET) treatment. The overall fertilization rate by ICSI is between 50% and 70% (1-3). In contrast to IVF, approximately 80% of unfertilized oocytes which have no signs of activation following ICSI have a spermatozoon in their ooplasm, and in the majority of these oocytes, the sperm head was partially or completely decondensed (4). Many studies have suggested that failure of oocyte activation is the principal cause of fertilization failure in ICSI (15-17). Even spermatozoa with abnormal morphology and elongated spermatids have an ability to activate oocytes similar to normal mature spermatozoa. Considering these findings, cell membranes of spermatozoa which fuse to oolemma during normal fertilization still remain in the ooplasm and may interfere with the migration of oscillin into ooplasm at the proper time, resulting in failure of oocyte activation. Recently, Yanagida *et al.*, reported that in 0.9% of ICSI cases with motile spermatozoa, the motile spermatozoa failed to fertilize the oocytes despite an apparently successful ICSI procedure (18). Furthermore, in patients with round-headed sperm the fertilization rate following ICSI was extremely low compared with patients




Fig.1. Model for the mechanisms of metaphase II arrest and release in vertebrate eggs. Fertilization triggers the activation of calmodulin-dependent kinase II via a Ca^{2+} transient, which probably activates the cyclin degradation machinery, resulting in cyclin B degradation and sister chromatid separation (Sagata, 1996)

with other types of male infertility due to the absence of oocyte activation capacity (19, 20). If unfertilized oocytes after ICSI are properly activated, they may form two pronuclei with extrusion of the second polar body, and cleave and develop normally. Indeed, oocytes injected with a spermatozoon unable to fertilize MII oocytes were immediately activated with electric pulse or calcium ionophore A23187. Consequently, successful pregnancies were achieved and two healthy babies were born (18, 20). The activation methods they used produced two pronuclei with extrusion of the second polar body in oocytes with a spermatozoon. This suggests that the activation method which produces one pronucleus with extrusion of the second polar body in oocytes without a spermatozoon is necessary for producing embryos with normal karyotypes.

OOCYTE ACTIVATION WITH CALCIUM IONOPHORE A23187 AND PUROMYCIN

There are many physical and chemical stimuli for inducing parthenogenetic or oocyte activation in mammals, such as ethanol (21, 22), calcium ionophore A23187 (Ca ionophore) (23, 24), inhibitors of protein synthesis (25, 26) and electric pulse (24, 27-29). Mouse aged oocytes are easily activated at a consistently high rate by Ca ionophore or ethanol which induce a calcium transient in the ooplasm (30). In contrast to mouse oocytes, Ca ionophore does not effectively activate human aged oocytes (31, 32). In general, human aged oocytes are less sensitive to widely used stimuli like Ca ionophore and ethanol than mouse aged oocytes. However, puromycin, an inhibitor of protein synthesis, produces parthenogenones in about 90% of human oocytes (31, 33, 34). However, 77% of the human parthenogenones activated by puromycin possessed 2 haploid pronuclei or 1 diploid pronucleus without extrusion of the second polar body (34). Moses and Masui showed that 6-dimethylaminopurine (6-DMAP)-an analogue of puromycin-suppressed extrusion of the second polar body in mouse oocytes (35). As mentioned above, an increase in intracellular Ca concentration and MPF inactivation are the main molecular events of M-G1 transition in MII oocytes and to mimic these events, human aged unfertilized oocytes following IVF were activated using various combinations of Ca ionophore and puromycin or 6-DMAP. When oocytes activated with 5 μ M calcium ionophore A23187 (Sigma, St.

Louis, Mo, USA) for 5 minutes were incubated in HTF medium containing 10 μ g/ml puromycin (Sigma, St. Louis, Mo, USA) for 5 hours, the activation rate was approximately 90%, which is similar to the rates reported by De Sutter et al. (33, 34) and Balakier et al. (31). Furthermore, in approximately 80% of the activated oocytes, one pronucleus was formed with extrusion of the second polar body (Fig 2). However, when oocytes were treated with Ca ionophore and 6-DMAP, they did not extrude the second polar body. In the next experiment, the activated oocytes displaying one pronucleus with extrusion of the second polar body were cytogenetically analysed. Cytogenetical analysis showed a normal haploid set of chromosomes in 78% of the activated oocytes. These findings suggested that the activated oocytes displaying one pronucleus with extrusion of the second polar body were haploid. If unfertilized oocytes with a spermatozoon are activated with a combination of Ca ionophore and puromycin, the activated oocytes should possess two pronuclei -one female and one male pronuclei- with extrusion of the second polar body similar to normal fertilized




Fig.2. The activation rate, proportion of parthenogenones displaying one pronucleus with extrusion of the second polar body and cleavage rate in human aged oocytes activated by calcium ionophore and protein phosphorylation inhibitors. Human aged unfertilized oocytes activated with 5 μ M calcium ionophore A23187 for 5 minutes were incubated in HTF medium containing 10 μ g/ml puromycin for 5 hours. After incubation for 13 hours, the oocytes were examined using an inverted microscope with Nomarski interference phase-contrast optics. Activated oocytes were defined as oocytes with at least one pronucleus or that had immediately cleaved.

*p<0.05 : puromycin and 6-DMAP vs control

**p<0.05 : puromycin vs 6-DMAP and control

***P<0.05 : 6-DMAP vs puromycin and control

oocytes. In our preliminary study, when unfertilized oocytes following ICSI were treated with Ca ionophore and puromycin, two pronuclei and the second polar body appeared in 30% of the oocytes 18 hours after treatment with Ca ionophore. Seventy percent of the activated oocytes developed as far as the 8-cell stage 42 hours after the Ca ionophore treatment and in cytogenetical analysis all cleaved parthenogenones (n=4) showed normal karyotypes. However, whether one of the appeared pronuclei was derived from a spermatozoa or not remains unknown. The sex ratio of the cleaved parthenogenones should be determined to clarify this, because half of the cleaved parthenogenones must have Y chromosomes if the genome derived from a spermatozoa is normally involved in the early development.

Young oocytes are not easily activated by a calcium transient. Kubiak *et al.* (22) and Vincent *et al.* (36) reported that mouse young oocytes activated with Ca ionophore or 7% ethanol did expel a second polar body, but became arrested again at so-called metaphase III (MIII), in which unichromatid chromosomes were scattered in misshapen microtubule spindles. However, 6-DMAP induced mouse young oocytes that were activated with Ca ionophore, but arrested at MIII, to enter interphase (35). In the present study, when mouse young oocytes were exposed to 5 μ M Ca ionophore and incubated in a medium containing 10 μ g/ml puromycin for 4 hours, the activation rate was 90%. More than 80% of the activated oocytes showed one pronucleus with extrusion of the second polar body (Table 1, Fig.3). These findings are similar to those of human aged oocytes. Furthermore, the DNA content of the pronucleus of the activated oocytes was half the DNA content of a set of MII chromosomes and the most highly activated oocytes possessed 20 chromosomes (normal haploid set). In conclusion, a

combination of Ca ionophore and puromycin produced haploid parthenogenones in young mouse oocytes. This strongly suggests that the combination is effective for inducing haploid parthenogenones from young human oocytes although we did not activate human young oocytes in the present study.

MOLECULAR EVENTS INDUCED BY CALCIUM IONOPHORE AND PUROMYCIN

It is unknown how the combination of Ca ionophore and puromycin produces parthenogenones displaying one pronucleus with extrusion of the second




Fig. 3. Parthenogenone displaying one pronucleus with extrusion of the second polar body. arrow : the second polar body, arrow head : pronucleus

Table 1. Effect of calcium ionophore and/or protein phosphorylation inhibitors on oocyte activation of mouse young oocytes.

		No. of oocytes treated	Activated oocytes (%)	
			Total No. of activated oocytes (%)	No. of parthenogenones with 1PN2PB (%)
	Control	90	8 (8.9)	3 (3.3)
Ca ionophore (+)	Puromycin	116	103* (88.8)	94* (81.0)
	DMAP	93	41 (44.1)	28 (30.1)

*p<0.0001 (the puromycin group vs the control and DMAP groups)

PN : pronucleus

PB : polar body

DMAP : 6-dimethylaminopurine

Activated oocytes means oocytes which had formed at least one pronucleus.

polar body. A Ca^{2+} transient induced by the Ca ionophore must trigger the M-G1 transition. Rinaudo *et al.* reported that human oocytes exposed to Ca ionophore exhibited an isolated elevation of intracellular calcium concentration, followed by a prompt return to baseline levels (32). A Ca^{2+} transient activates calmodulin-dependent kinase II (CaM KII), which probably turns on the cyclin degradation machinery, resulting in cyclin B degradation and sister chromatid separation (9). In the present study a Ca^{2+} transient induced by Ca ionophore probably inactivated MPF activity via CaM KII. However, a Ca^{2+} wave could be a positive signal that activates myosin light chain kinase (37). This kinase is activated through the Ca^{2+} /calmodulin complex and activates the contractile ring which consists of actin and myosin in cytokinesis. The extrusion of the second polar body is an uneven cleavage of the cell. A Ca^{2+} transient induced by Ca ionophore might play an important role of extruding the second polar body.

In our experiments, 25-50% of oocytes activated by Ca ionophore alone extruded the second polar body without formation of a pronucleus (M III stage). Cotreatment with puromycin increased the rate of oocytes with a pronucleus or pronuclei to approximately 90%. Verlhac *et al.* reported that when freshly ovulated MII mouse oocytes were treated with ethanol, MPF activity dropped to one fourth at the time of polar body extrusion and this activity increased again in MIII oocytes to a level slightly lower than that observed in MII oocytes (38). MPF activity of MIII oocytes might have increased again in the present study. Puromycin and 6-DMAP are known to inhibit mitogen activating protein kinase (MAPK) which is a component of CSF in mouse (39) and bovine oocytes (40). Sagata reported that MAPK probably activates a positive regulator of MPF (e.g. cdc 25) or inhibits negative regulators (e.g. weel or proteinphosphatase 2 A) (9). So, this suggests that puromycin decreased MPF activity via suppression of MAPK, resulting in the formation of a pronucleus.

Cotreatment with 6-DMAP completely suppressed extrusion of the second polar body in human aged oocytes, but not cotreatment with puromycin. This finding agrees with the study by Liu *et al.* (41) that the combination of Ca ionophore and 6-DMAP disrupted the oocyte spindle and effectively produced parthenogenones displaying one pronucleus without extrusion of the second polar body. However, treatment of puromycin without Ca ionophore produces

parthenogenones displaying 2 haploid pronuclei or 1 diploid pronucleus without extrusion of the second polar body. It is unknown how the combination of Ca ionophore and puromycin can extrude the second polar body. Some molecular events provoked by the Ca ionophore might be suppressed by 6-DMAP, but not puromycin, resulting in suppression of extrusion of the second polar body.

A major difference between puromycin and 6-DMAP is that puromycin inhibits protein synthesis, while 6-DMAP does not. Rime *et al.* (42) reported that puromycin provokes a approximate 60% decrease in protein synthesis together with the formation of a pronucleus. However, the emission of the second polar body does not appear to be attributed to the inhibition of protein synthesis.

CLINICAL ADAPTATION OF THE OOCYTE ACTIVATION

Clinical adaptation of this method may be possible, but it is still far from reality at present. Prior to the clinical adaptation, the safety of rescuing unfertilized oocytes and the cytotoxic, teratogenic and mutagenetic activity of Ca ionophore and puromycin should be approved. Many cytogenetical analyses show that approximately 25% of unfertilized oocytes following IVF have chromosomal aberrations (11, 14, 15, 43, 44, 45-47). Pickering *et al.* reported that aged unfertilized oocytes (24-48 h) displayed a higher incidence (approximately 30%) of disrupted or abnormal spindles (48). Furthermore, Dozortsev recently reported that swollen nuclei with fragmented pieces of chromosomes were observed in 51% of unfertilized oocytes after ICSI and these probably corresponded to degenerating nuclei (16). If unfertilized oocytes are evenly activated with Ca ionophore and puromycin, activated oocytes with chromosomal aberrations and/or swollen nuclei with fragmented pieces of chromosomes must be discarded. However, in the present study approximately 80% of oocytes, which had been unfertilized after IVF, showed a normal haploid set of chromosomes after activation. This may suggest that MII oocytes responsive to Ca ionophore and puromycin have 23 chromosomes with 2C and have an ability to finish meiosis. In respect to organelles other than chromosomes, Sousa and Tesarik reported that no obvious ultrastructural signs of degeneration (vacuolation, mitochondrial swelling) were apparent in any of 15 oocytes that failed to fertilize by 44 h

after ICSI although the functions of organelles in the oocytes remain unknown (49).

In ICSI cases with spermatozoa unable to fertilize oocytes, successful pregnancies were achieved using a combination of ICSI and oocyte activation (18, 19). Young oocytes are difficult to activate because the activities of MPF and MAPK are stable. However, in mice, 90% of newly ovulated oocytes were activated with Ca ionophore and puromycin and approximately 80% of the activated oocytes showed one pronucleus with extrusion of the second polar body. This suggests that the combination of Ca ionophore and puromycin is effective for activating young human oocytes. In cases of activating young oocytes, the adverse effects of aging on oocyte quality was not found to be an important consideration.

Ca ionophore A23187 does not appear to be cytotoxic to oocytes if the concentration is low (50). Indeed, in mice, oocytes activated by Ca ionophore developed as far as the post-implantation stage. If oocytes survive an isolated elevation of intracellular calcium concentration induced by Ca ionophore, the activated oocytes with a diploid set of chromosomes developed as far as the post-implantation stage. However, these parthenogenones do not have a paternal genome and some genes like the *igf2* gene are not expressed because of genomic imprinting. So, the ability to develop as far as the post-implantation stages does not mean that the genes are normal. Further studies are required to approve the safety of Ca ionophore, although a healthy baby was born after treatment with Ca ionophore.

A high dosage and/or prolonged treatment with puromycin can irreversibly damage the chromosomes (pulverization, breakage), but not a low dosage and a short treatment (33). Indeed, in the present human and mouse cytogenetical analysis, 70–80% of oocytes activated with Ca ionophore and puromycin showed a normal haploid set of chromosomes with normal morphology. Bovine oocytes activated with Ca ionophore and 6-DMAP developed to the blastocyst and hatched blastocyst stages similar to those derived from IVF-ET (41). These findings suggest that inhibition of MAPK activity do not appear to affect the developmental ability of embryos.

At present there are no methods to select fresh normal embryos. By culturing embryos to the blastocyst stage, genetically abnormal embryos may be selected out through aberrations in their development (51) and may produce the most viable

embryos without abnormal organelles. However, the ability to develop as far as the post-implantation stages does not indicate the genes are developing normally. At present, the teratogenetic and mutagenetic activity of Ca ionophore and puromycin are under investigation in our center using a mouse model.

REFERENCES

1. Palermo G, Joris H, Derde MP, Camus M, Devroey P, Van Steirteghem AC : Sperm characteristics and outcome of human assisted fertilization by subzonal insemination and intracytoplasmic sperm injection. *Ferti Steril* 59 : 826-835, 1993
2. Van Steirteghem AC, Nagy Z, Joris H, Liu J, Staessen C, Smits J, Wisanto A, Devroey P : High fertilization and implantation rates after intracytoplasmic sperm injection. *Hum Reprod* 8 : 1061-1066, 1993
3. Payne D, Matthews CD : Intracytoplasmic sperm injection-clinical results from the Reproductive Medicine Unit, Adelaide. *Reprod Fertil Dev* 7 : 219-227, 1995
4. Flaherty SP, Payne D, Matthews CD : Fertilization failures and abnormal fertilization after intracytoplasmic sperm injection. *Hum Reprod* 13 (suppl 1) : 155-164, 1998
5. Sagata N, Watanabe N, George F, Vande W, Ikawa Y : The *c-mos* proto-oncogene product is a cytostatic factor responsible for meiotic arrest in vertebrate eggs. *Nature* 342 : 512-518, 1989
6. Homa ST, Carroll J, Swann K : The role of calcium in mammalian oocyte maturation and egg activation. *Hum Reprod* 8 : 1274-1281, 1993
7. Taylor CT, Lawrence YM, Kingsland R, Biljan MM, Cuthbertson KS : Oscillations in intracellular free calcium induced by spermatozoa in human oocytes at fertilization. *Hum Reprod* 8 : 2174-2179, 1993
8. Swann K, Ozil JP : Dynamics of the calcium signal that triggers mammalian egg activation. *Int Rev Cytol* 152 : 183-222, 1994
9. Sagata N : Meiotic metaphase arrest in animal oocytes : its mechanisms and biological significance. *Trend Cell Biol* 6 : 22-28, 1996
10. Schmiady H, Kentenich H : Premature chromosome condensation after in-vitro fertilization.

- Hum Reprod 4 : 689-695, 1989
11. Ma S, Kalousek D, Zouves C, Yuen BH, Gomel V, Moon YS : Chromosome analysis of human oocytes failing to fertilize in vitro. *Fertil Steril* 51 : 992-997, 1989
 12. Balakier H, Casper RF : A morphologic study of unfertilized oocytes and abnormal embryos in human in vitro fertilization. *J Assist Reprod Genet* 8 : 73-79, 1991
 13. Van Blerkom J, Davis P, Merriam J : A retrospective analysis of unfertilized and presumed parthenogenetically activated human oocytes demonstrates a high frequency of sperm penetration. *Hum Reprod* 9 : 2381-2388, 1994
 14. Wall MB, Marks K, Smith TA, Gearon CM, Muggelton-Harris AL : Cytogenetic and Fluorescent in-situ hybridization chromosomal studies on in-vitro fertilized and intracytoplasmic sperm injected failed-fertilized human oocytes. *Hum Reprod* 11 : 2230-2238, 1996
 15. Edirisinghe WR, Murch A, Junk S, Yovich JL : Cytogenetic abnormalities of unfertilized oocytes generated from in-vitro fertilization and intracytoplasmic sperm injection : a double-blind study. *Hum Reprod* 12 : 2784-2791, 1997
 16. Dozortsev D, De Sutter P, Dhont M : Behavior of spermatozoa in human oocytes displaying no or one pronucleus after intracytoplasmic sperm injection. *Hum Reprod* 9 : 2139-2144, 1994
 17. Sousa M, Tesarik J : Ultrastructural analysis of fertilization failure after intracytoplasmic sperm injection. *Hum Reprod* 9 : 2374-2380, 1994
 18. Yanagida K, Katayose H, Yazawa H, Kimura Y, Sato A, Yanagimachi H, Yanagimachi R : Successful fertilization and pregnancy following ICSI and electrical oocyte activation. *Hum Reprod* 14 : 1307-1311, 1999
 19. Rybouchkin A, De Sutter P, Straeten FV, Dhont M, Quatacker J : Fertilization and pregnancy after assisted oocyte activation and intracytoplasmic sperm injection in a case of round-headed sperm associated with deficient oocyte activation capacity. *Fertil Steril* 68 : 1144-1147, 1997
 20. Battaglia DE, Koehler JK, Klein NA, Toker MJ : Failure of oocyte activation after intracytoplasmic sperm injection using round-headed sperm. *Fertil Steril* 68 : 118-122, 1997
 21. Nagai T : Parthenogenetic activation of cattle follicular oocytes in vitro maturation and with ethanol. *Gamete Res* 16 : 143-249, 1987
 22. Kubiak JZ : Mouse oocytes gradually develop the capacity for activation during the metaphase II arrest. *Dev Biol* 136 : 537-545, 1989
 23. Meyerhof PG, Masui Y : Ca and Mg control of cytostatic factors from Rana pipiens oocytes which cause metaphase and cleavage arrest. *Dev Biol* 61 : 214-229, 1977
 24. Ware CB, Barnes FL, Maiki-Lauria M, First NL : Age dependence of bovine oocyte activation. *Gamete Res* 22 : 265-275, 1989
 25. Fulka JJr, Leibfried-Rutledge ML, First NL : Effect of 6-dimethylaminopurine on germinal vesicle breakdown of bovine oocytes. *Mol Reprod Dev* 29 : 379-384, 1991
 26. Siracusa G, Whittingham DG, Molinaro M, Vivarelli E : Parthenogenetic activation of mouse oocytes induced by inhibitors of protein synthesis. *J Embryol Exp Morphol* 43 : 157-166, 1978
 27. Powell R, Barnes FL : The kinetics of oocyte activation and polar body formation in bovine embryo clones. *Mol Reprod Dev* 33 : 53-58, 1992
 28. Campbell KHS, Ritchie WA, Wilmut I : Nuclear-cytoplasmic interactions during the first cell cycle of nuclear transfer reconstructed bovine embryo : Implications for deoxyribonucleic acid replication and development. *Biol Reprod* 49 : 933-942, 1993
 29. Prochazka R, Durnford R, Fiser PS, Marcus GJ : Parthenogenetic development of activated in vitro matured bovine oocytes. *Theriogenology* 39 : 1025-1032, 1993
 30. Cuthbertson KSR, Whittingham DG, Cobbold PH : Free Ca²⁺ increase in exponential phases during mouse oocyte activation. *Nature* 294 : 754-757, 1981
 31. Balakier H, Casper RF : Experimentally induced parthenogenetic activation of human oocytes. *Hum Reprod* 8 : 740-743, 1993
 32. Rinaudo P, Pepperll JR, Buradgunta S, Massobrio M, Keef DL : Dissociation between intracellular calcium elevation and development of human oocytes treated with calcium ionophore. *Fertil Steril* 68 : 1086-1092, 1997
 33. De Sutter P, Dozortsev D, Cieslak J, Wolf G, Verlinsky Y, Dyban A : Parthenogenetic activation of human oocytes by puromycin. *J Assist Reprod Genet* 9 : 328-337, 1992
 34. De Sutter P, Dozortsev D, Wolf G, Vrijens P, Desmet R, Dhont M : Cytogenetic analysis of human oocytes parthenogenetically activated by puromycin. *J Assist Reprod Genet* 11 : 382-

- 388, 1994
35. Moses RM, Masui Y ; Enhancement of mouse egg activation by the kinase inhibitor, 6-dimethylaminopurine (6-DMAP). *J Exp Zool* 270 : 211-218, 1994
 36. Vincent C, Cheek TR, Johnson MH : Cell cycle progression of parthenogenetically activated mouse oocytes to interphase is dependent on the level of internal calcium. *J Cell Sci* 130 : 389-396, 1992
 37. Satterwhite LL, Pollard TD : Cytokinesis. *Current Biol* 4 : 43-52, 1992
 38. Verlhac MH, Kubiak JZ, Clarke HJ, Maro B : Microtubule and chromatin behavior follow MAP kinase activity but not MPF activity during meiosis in mouse oocytes. *Development* 120 : 1017-1025, 1994
 39. Verlhac MH, De Pennart H, Maro B, Cobb MH, Clarke H : MAP kinase becomes stably activated at metaphase and is associated with microtubule-organizing centers during meiotic maturation of mouse oocytes. *Dev Biol* 158 : 330-340, 1993
 40. Liu L, Ju JC, Yang X : Differential inactivation of maturation-promoting factor and mitogen-activated protein kinase following parthenogenetic activation of bovine oocytes. *Biol Reprod* 59 : 537-545, 1998
 41. Liu L, Ju JC, Yang X : Parthenogenetic development and protein patterns of newly matured bovine oocytes after chemical activation. *Mol Reprod Dev* 49 : 298-307, 1998
 42. Rime H, Neant I, Guerrier P, Ozon R : 6-dimethylaminopurine(6-DMAP), a reversible inhibitor of the transition to metaphase during the first meiotic cell division of the mouse oocyte. *Dev Biol* 133 : 169-179, 1989
 43. Martin RH, Mahadevan MM, Taylor PJ, Hildebrand K, Long-Simpson L, Peterson D, Yamamoto J, Fleetham J : Chromosomal analysis of unfertilized human oocytes. *J Reprod Fertil* 78 : 673, 1986
 44. Wramsby H, Fredga K : Chromosome analysis of human oocytes recovered from preovulatory follicles in stimulated cycles. *N Engl J Med* 316 : 121, 1987
 45. Pellestor F, Sele B : Assessment of aneuploidy in human female by using cytogenetics of IVF failures. *Am J Hum Genet* 42 : 274, 1988
 46. Plahot M, Veigu A, Montagut J, Grouchy J, Calderon G, Lepretre S, Junka A, Santalo J, Elisabeth C, Mandelbaum J, Barri P, Degony J, Cohen J, Egozcue J, Sabatier JC, Salat-Baroux J : Are clinical and biological IVF parameters correlated with chromosomal disorders in early life : a multicentric study. *Hum Reprod* 3 : 627, 1988
 47. Bonso A, Chye NS, Ratnam S, Sathanantan H, Wong PC : Chromosome anomalies in human oocytes failing to fertilize after insemination in vitro. *Hum Reprod* 3 : 645, 1988
 48. Pickering SJ, Johnson MH, Braude PR, Houlston E : Cytoskeletal organization in fresh, aged and spontaneously activated human oocytes. *Hum Reprod* 8 : 978-989, 1988
 49. Sousa M, Tesarik J : Ultrastructural analysis of fertilization failure after intracytoplasmic sperm injection. *Hum Reprod* 9 : 2374-2380, 1994
 50. Tesarik J, Sousa M : More than 90% fertilization rates after intracytoplasmic sperm injection and artificial induction of oocyte activation with calcium ionophore. *Fertil Steril* 63 : 343-349, 1995
 51. Bongso A : Why blastocyst culture? *Handbook on blastocyst culture*. Sydney Press Indusprint, Singapore, 1999